

(to appear in the Astronomical Journal)

The Universality of the Fundamental Plane of E and S0 Galaxies. Sample Definition and I Band Photometric Data

Marco Scodeggio^{1,2}, Riccardo Giovanelli¹ and Martha P. Haynes

Center for Radiophysics and Space Research and National Astronomy and Ionosphere Center³,
Cornell University, Ithaca, NY 14853

mscodegg@eso.org

riccardo, haynes@astrosun.tn.cornell.edu

ABSTRACT

As part of a project to compare the Fundamental Plane and Tully-Fisher distance-scales, we present here I band CCD photometry for 636 early-type galaxies in 8 clusters and groups of galaxies. These are the A262, A1367, Coma (A1656), A2634, Cancer and Pegasus clusters, and the NGC 383 and NGC 507 groups. Sample selection, cluster properties, and cluster membership assignment criteria are discussed. We present photometric parameters that are used in the Fundamental Plane relation, the effective radius r_e and the effective surface brightness μ_e , as derived from a $r^{1/4}$ fit to the observed radial photometric profile of each galaxy. A comparison with similar data found in the literature for the Coma cluster shows that large systematic uncertainties can be introduced in the measurement of r_e and μ_e by the particular method used to derive those parameters. However the particular combination of these two parameters that enters in the Fundamental Plane relation is a quantity that can be measured with high accuracy.

Subject headings: distance scale — galaxies: distances and redshifts — galaxies: elliptical and lenticular — galaxies: fundamental parameters — galaxies: photometry

¹Visiting Astronomer, Kitt Peak National Observatory, National Optical Astronomy Observatories, which is operated by the Association of Universities for Research in Astronomy, Inc. (AURA) under cooperative agreement with the National Science Foundation.

²Present address: European Southern Observatory, Karl-Schwarzschild-Straße 2, D-85748, Garching bei München, Germany.

³The National Astronomy and Ionosphere Center is operated by Cornell University under a cooperative agreement with the National Science Foundation.

1. Introduction

This is the third in a series of papers (Scodeggio, Giovanelli & Haynes 1997a,b; hereafter Paper I and Paper II) devoted to a study of the universality of the Fundamental Plane (FP) relation of early-type galaxies (Djorgovski & Davis 1987; Dressler *et al.* 1987). This relation, and the equivalent Tully-Fisher (TF) relation for spiral galaxies (Tully & Fisher 1977), have been widely used in the recent past to produce large samples of redshift-independent distance estimates. The small scatter of the FP relation has been also used to put constraints on the epoch of early-type galaxy formation, and on the subsequent evolution and star formation histories of those objects (see for example Guzmán, Lucey & Bower 1993; Renzini & Ciotti 1993; van Dokkum & Franx 1996; Pahre & Djorgovski 1997).

Both applications are based on the assumption that the FP and TF relations are universal: all galaxies in all locations and environments have effective radius, effective surface brightness and velocity dispersion, or magnitude and rotation velocity that are related by the same relation, within measurement uncertainties and the intrinsic dispersion present in the relations. However, this might not be the case, since a number of environmental effects are known to exist that have the potential of altering significantly the stellar population and star formation history of the affected galaxies. Therefore, the FP and TF universality should not be taken for granted, but should rather be observationally verified (discussions on this subject are presented by Djorgovski, de Carvalho & Han 1988; Guzmán *et al.* 1992; Jørgensen, Franx & Kjaergaard 1996; Schroeder 1996; Giovanelli *et al.* 1997b; and in Paper I).

The most direct way to test the reliability of a distance indicator relation is to compare its predictions with those of another independent, and at least equally reliable, method. The TF and FP relations provide the opportunity for such a comparison. They have similar accuracy, and they are independent methods, because they are applied to galaxies of different morphological type and different clustering properties. Because of the limited accuracy of the distance estimates obtained with the TF and FP relation, a direct comparison of the two methods can be done most effectively using clusters of galaxies, where large samples can be used to reduce the statistical uncertainty in the estimate of the cluster distances. It should be kept in mind, however, that given the existing correlation between cluster richness and the properties of the intracluster medium (see for example Edge & Stewart 1991), it is desirable that such a comparison bridge a set of clusters characterized by a wide range of richness, in order for the TF-FP comparison to have broad validity.

In a recent work, Giovanelli *et al.* (1997a,b; hereafter G97a,b) have presented TF measurements for a large sample of cluster objects. This represent the largest dataset currently available that can be used for estimating cluster distances using the TF relation. In this and the following papers of this series, we present the results of a FP study of a subset (eight clusters) of the clusters studied by G97a,b. The main purpose of this work is to obtain distance estimates for those clusters, with accuracies comparable to those achievable using the TF dataset, to be used for a comparison between the two methods. In this paper, we briefly describe the cluster sample selection and the

adopted cluster membership criteria, and we present I band CCD photometry for 636 early-type galaxies in those clusters. In a companion paper (Scodeggio, Giovanelli & Haynes 1998a), we present the results of spectroscopic observations of a subset of these galaxies. Following that work, we will discuss the properties of the I band FP relation, and the comparison between TF and FP distance estimates for the clusters in our sample (Scodeggio, Giovanelli & Haynes 1998b).

The cluster sample selection is described in Section 2; observations and the data reduction procedure are described in Section 3 and 4; the photometric sample is presented in Section 5. An assesment of the quality of the photometric data presented in this work is presented in Section 6. Throughout the paper we parametrize distance–dependent quantities via $H_0 = 100h$ $\text{km s}^{-1}\text{Mpc}^{-1}$.

2. Cluster sample selection and membership

Recently G97a,b have presented TF measurements for a sample of $\simeq 800$ galaxies in 24 clusters. Because one of the goals of that work was to obtain a kinematical rest frame that could be used to study the local peculiar velocity field, it was important that such sample of clusters had a rather uniform sky coverage, so that localized anomalies in the velocity field would average out in the global sample. On the contrary, the main goal of this work is a detailed, point-by-point comparison of distance estimates, irrespective of the possible peculiar velocities that the target clusters might have. Therefore a uniform sky coverage is not an important requirement for our sample. It is essential instead to cover a wide spread in cluster richness and the properties of the intracluster medium (ICM), since known environmental effects on galaxies are strongly dependent on the local galaxy density, or on the density of the ICM. We have therefore selected a subset of the clusters in G97a,b as targets for the FP study, restricted to the northern hemisphere, mainly to facilitate the spectroscopic observations. The set is composed of the 8 clusters and rich groups of galaxies NGC 383 group, NGC 507 group, A262, Cancer, A1367, Coma (A1656), Pegasus, A2634.

2.1. Cluster properties

Table 1 lists the most important properties of these clusters and groups. The cluster center is generally assumed to be coincident with the position of the brightest galaxy, except in the case of the irregular cluster A1367, where the center is assumed to be at the mid-point between the two brightest ellipticals. Systemic heliocentric velocity and velocity dispersion are taken from the literature, according to the references quoted in the table. Heliocentric velocities are converted to the cosmic microwave background (CMB) reference frame using the CMB dipole determination from Kogut *et al.* (1993). Population fractions data are from Dressler (1980), and Giovanelli & Haynes (1985), while the fraction of HI deficient objects in a cluster (this is the fraction of spiral galaxies that have HI deficiency > 0.3 , according to the definition of Giovanelli & Haynes 1985

and Haynes & Giovanelli 1984) is taken from Giovanelli & Haynes (1985), and Magri *et al.* (1988). X-ray data are taken from David *et al.* (1993).

The most extreme properties within our sample are those of the Coma cluster, a rich, dynamically relaxed, and X-ray luminous cluster, that exhibits strong morphological segregation and HI deficiency among the spiral galaxy population. Then we have medium richness clusters like A2634 and A1367: the first is a dynamically evolved cluster (see however Pinkney *et al.* 1993 and Schindler & Prieto 1997 for a discussion on a possible recent merger event in A2634), exhibiting clear morphological segregation and pronounced HI deficiency among the spiral galaxy population, while the second is an irregular cluster, with only moderate morphological segregation and HI deficiency. A262 is a transitional cluster, where low X-ray luminosity and velocity dispersion, cool ICM temperature, and high spiral fraction, all indicative of a dynamically unevolved state, coexist with the presence of a massive dominant galaxy, which is typical of more evolved systems (see for example Forman & Jones 1982, and Sarazin 1988). The cluster exhibits moderate morphological segregation and HI deficiency. The low end of the cluster richness distribution is represented by the Pegasus and Cancer clusters. Both are spiral rich, low X-ray luminosity objects, that exhibit little morphological segregation and marginal HI deficiency. Here we refer to Cancer as the largest of the separate groups that were identified by Bothun *et al.* (1983) over the area and redshift range traditionally associated with the Cancer cluster. This is group A in the nomenclature of Bothun *et al.* The rich groups associated with the galaxies NGC 507 and NGC 383 are often considered as a single entity, the so-called Pisces cluster. However Sakai, Giovanelli & Wegner (1994) have clearly shown that the two can be separated. In both groups the galaxy population is dominated by elliptical and S0 galaxies, and HI deficiency is almost absent among the spiral galaxy population.

2.2. Cluster membership

A discussion of the membership criteria adopted for the TF work on the clusters in the present sample was presented by G97a (their section 8). Here we adopt the same basic criteria, with the advantage that the bulk of the early-type galaxy population is concentrated towards the cluster centers, where the application of those criteria is easier than at the cluster periphery. Therefore we do not separate our FP sample into certain and possible cluster members, as done in G97a,b, but we define only *bona-fide* cluster members. The photometric sample presented here, however, includes also a number of galaxies for which spectroscopic data is not available, considered likely cluster members on the basis of their size and I band flux.

Whenever caustic lines in redshift space (Kaiser 1987, Regös & Geller 1989, van Haarlem 1992) are available for one of the clusters in our sample, membership determination is based on those lines, as estimated for $\Omega_0 = 0.3$. This is the case for Coma, where caustic lines have been published by Regös & Geller (1989), van Haarlem (1992), and Gavazzi, Randone & Branchini (1995); A2634, where they have been published by Scodeggio *et al.* (1995); and A1367, where they have been published by Regös & Geller (1989), and Gavazzi *et al.* (1995). For Coma and A1367

we use the caustic lines published by Regös & Geller (1989).

For A262, we use the detailed analysis of the spatial and velocity distribution of the galaxies in the A262 area presented by Sakai *et al.* (1994). Although they did not derive redshift space caustic lines for this cluster, their work can be used to derive reliable membership criteria, out to a radial distance of 2° from the cluster center, beyond which the contamination from the sparse population of the Perseus-Pisces Supercluster becomes significant. We define as A262 members all galaxies within 2° from the cluster center, with cz_{hel} within the interval $3500\text{--}6300\text{ km s}^{-1}$. Pegasus is well separated in redshift space from foreground and background objects, and membership assignments are not problematic, especially for the early-type population. We define as Pegasus members all galaxies within 2° from the cluster center, with cz_{hel} within the interval $2700\text{--}5100\text{ km s}^{-1}$. For Cancer, we use the membership table published by Bothun *et al.* (1983) for group A. For the NGC 383 and NGC 507 groups, we use the analysis published by Sakai *et al.* (1994). Although redshift space caustic lines could not be reliably derived, because of the small samples involved, membership can be assigned confidently, thanks to the relative under-density of the region in front and behind the two groups. Here we define as NGC 507 members all galaxies within 1.5° from the group center, with cz_{hel} within the interval $3500\text{--}6700\text{ km s}^{-1}$, and as NGC 383 members all galaxies within 1.0° from the group center, with cz_{hel} within the interval $3800\text{--}6500\text{ km s}^{-1}$.

3. Observations

All photometric measurement have been derived from I-band CCD images. The adoption of I band for photometry was mainly dictated by the practical notion that the cluster images contain several galaxies per frame, both of early- and late-type, and thus can be useful for TF application, where the I band is favored for compatibility with G97a,b.

Most of the observations were obtained during three runs with the 0.9m telescope of the Kitt Peak National Observatory (KPNO). A small number of observations in the cluster A2634 was obtained during observing runs with the 1.3m telescope of the Michigan-Dartmouth-MIT (MDM) Observatory (runs 1 and 3 in Table 2). Other observations in A2634 were obtained by KPNO staff with the 0.9m telescope during a period of service observations (run 2 in Table 2). Table 2 lists all observing runs during which data were obtained for this project, and gives parameters of the instrumental setups. The most significant difference between the MDM and the KPNO observations is the much smaller field of view of the 1.3m MDM telescope, when compared with the 0.9m KPNO telescope.

All CCD frames were obtained with 600 seconds integration time. For the brightest objects, a second frame with shorter integration (200–300 seconds) was necessary to avoid saturation of the image in the nuclear region of the galaxy. The median effective seeing (measured as the FWHM of the stellar light profile) for the entire data set is $1.6''$. Figure 1 shows the distribution of effective seeing for the galaxy observations. The actual atmospheric seeing might have been smaller at

times, but the pixel size with which the majority of the observations were obtained limits the effective seeing to $\geq 1.2''$. Given the large field of view of the KPNO 0.9m telescope, and the large density of galaxies in the cluster environments studied here, a complete mapping of the clusters' central regions was obtained, with a mosaic of frames on a regularly spaced grid. Only for the galaxies at the periphery of the various clusters off-grid pointings were used.

4. Data reduction

4.1. Basic reduction

The data reduction process was performed entirely using a combination of standard and specially written IRAF⁴ procedures. All frames were bias-subtracted, and flat-fielded using a “superflat” obtained from a median-filtered combination of a large ($n > 40$) number of frames. At this point, for ease of manipulation, one small frame (512 x 512 pixels) for each target galaxy was extracted from the 2048 x 2048 KPNO pixel frames. The 1024 x 1024 pixel MDM frames were kept as single frames. The sky background was subtracted from all frames, using the average number of counts measured in 10–12 “empty-sky” regions. The typical number of pixels over which this average was computed is 30,000. The accuracy of the background determination can be estimated from the comparison of the independent values subtracted from different 512 x 512 pixel frames extracted from the same 2048 x 2048 pixel frame. This uncertainty is typically of 0.2%. All pixels contaminated by the light of foreground stars or other galaxies, or by cosmic rays hits were blanked and excluded from the final steps of surface photometry.

For galaxies severely contaminated by the light of a nearby object (either another galaxy or a foreground star), an iterative subtraction procedure was used to obtain the individual light profiles. First, a model was fitted to the brighter object, after blanking the fainter one(s). After subtracting such model from the original image, a model was fitted to the fainter object, which again was subtracted from the original image. The procedure was iteratively applied until convergence was reached, usually at the second iteration, producing companion-subtracted light profiles, although the uncertainty on the photometry of these objects is somewhat higher than for isolated galaxies. This approach is especially necessary in the inner parts of clusters, where the galaxy density is high.

4.2. Photometric calibration

One or two Landolt (1992) fields, containing each one from 5 to 7 stars, were repeatedly observed during each night, at airmasses between 1.2 and 2.5, to provide estimates of the

⁴IRAF (Image Reduction and Analysis Facility) is distributed by the National Optical Astronomy Observatories.

photometric zero-point and atmospheric extinction coefficient. Observations at I and R band were used to obtain color corrected photometric solutions. Typical uncertainties in the zero-point determination are 0.02 mag, both at I and R band. All galaxy magnitudes were computed assuming a constant R–I color of 0.70 for elliptical, S0, and S0a galaxies (see, for example, Fukugita, Shimasaku & Ichikawa 1995). The uncertainty introduced by this assumption is negligible, because the color term coefficient is always very small (< 0.05). Galaxy observations were all obtained at airmass < 1.3 , thereby minimizing the effects of atmospheric extinction.

Most observations were obtained in photometric conditions. Those that were not were calibrated using either short exposure frames obtained in photometric conditions (run 2 was calibrated entirely in this way, using frames obtained during run 3), or taking advantage of the partial overlap among frames in the mapping grid (non photometric nights in runs 4, 5, 7, 8 were calibrated using this technique). In either case 10–15 stars that are present in both the “photometric” and the “non photometric” image were selected, and instrumental (i.e. non calibrated) magnitudes for all them were obtained in both images. After rejecting the obvious variable stars, mean magnitude differences between the stars in the two images were used to calibrate the “non photometric” frames. The typical uncertainty of this process is $\simeq 0.03$ mag.

4.3. Surface photometry

Surface photometry was obtained using the GALPHOT software package, written for IRAF/STSDAS⁵ mainly by W. Freudling, J. Salzer and M. Haynes. A galaxy’s light distribution was fitted with elliptical isophotes, using a modified version of the STSDAS *Isophote* package. The center, ellipticity and position angle of the ellipse were free parameters, while the ellipse semi-major axis was incremented by a fixed fraction of its value at each step of the fitting procedure. The procedure was stopped when the fitted mean number of counts per pixel dropped below the rms dispersion around the mean value. For this dataset, this typically happens close to the 23.5 mag arcsec^{−2} isophote. The fitted parameters yielded a model of the galaxy, which was used to compute integrated magnitudes as a function of semi-major axis. At this point the measured fluxes in counts per pixel were transformed into magnitudes per square arcsec, and all magnitudes were calibrated, color-corrected, and corrected for atmospheric extinction.

The radial surface brightness profiles were fitted with a $r^{1/4}$ profile (de Vaucouleurs 1948), to determine the effective radius r_e (the radius that contains half of the galaxy total light) and the effective surface brightness μ_e (the mean surface brightness within the effective radius). For all galaxies, the fit was performed from an inner radius twice as large as the seeing disk radius of the relative image. The outer radius was chosen depending on the shape of the surface brightness

⁵STSDAS (Space Telescope Science Data Analysis System) is distributed by the Space Telescope Science Institute, which is operated by AURA, under contract to the National Aeronautics and Space Administration.

profile. For those objects that are well described by the $r^{1/4}$ law at all radii, the fit was extended to the outermost isophotes. For all other galaxies, only the central core was fitted. The effective radius r_e and the surface brightness at r_e were the two free parameters in the fit, and the effective surface brightness μ_e was obtained from $\mu(r_e)$ as $\mu_e = \mu(r_e) - 1.392$ (this is strictly valid only for a galaxy that follows the $r^{1/4}$ profile at all radii). The median value for the uncertainty of the fits is 6.2% in r_e , and 0.07 mag arcsec $^{-2}$ in μ_e .

Integrated fluxes within the faintest reliably measured isophote were obtained. Total magnitudes were derived from those, by adding a term which was obtained by extrapolating the surface brightness profile to infinity. For those galaxies not well represented by the $r^{1/4}$ law at all radii the outer portion of the profile was fitted using either a different $r^{1/4}$ profile from the one used to fit the core of the galaxy, or an exponential disk profile, depending on the profile shape. The median uncertainty with which total magnitudes were derived is 0.06 mag.

Total magnitudes and effective surface brightnesses were corrected for Galactic extinction and cosmological effects. Extinction corrections were computed for each galaxy following the prescriptions of Burstein & Heiles (1978), using $A_I = 0.45A_V$. These corrections range from 0.0 to 0.13 mag. Statistically equivalent results are obtained using the extinction maps recently published by Schlegel, Finkbeiner & Davis (1998). Cosmological K-corrections were computed assuming $F_\nu \sim \nu^0$ (the energy distribution of early-type galaxies in the red is nearly flat) so that $K_I = 2.5 \log(1 + z)$, and range from 0.01 to 0.04 mag. The $(1+z)^4$ cosmological corrections to surface brightness range from 0.04 to 0.15 mag arcsec $^{-2}$. Effective radius and effective surface brightness were corrected for the smearing effects of seeing, following the prescriptions of Saglia *et al.* (1993a, see in particular their Figure 8). The seeing correction produced a median reduction of 2% of the measured values of r_e , and a median correction of 0.016 mag arcsec $^{-2}$ to the measured values of μ_e .

5. The photometric sample

In each cluster, surface photometry was obtained for E, S0, and S0a galaxies, potentially good candidates for FP work. A few galaxies that were classified as S0a/Sa or Sa were included in the sample because a velocity dispersion measurement was already available for them. Known foreground or background objects were excluded, while a number of galaxies without redshift measurement, but with flux and size that make them likely cluster members, were instead included in the samples, except in the case of the Cancer cluster, where foreground and background contamination is too pronounced. Galaxies in close pairs or multiplets, and galaxies contaminated by bright foreground stars projected over the galaxy or in its immediate vicinity were rejected, unless a velocity dispersion measurement was already available. Only in these cases was an attempt made to subtract the contribution of the contaminating object to the observed light distribution (see section 4.1). In total, photometric properties for 636 galaxies were measured. The final samples are close to magnitude-limited in nature, as will be seen in the discussion on the

FP relation (Scodeggio *et al.* 1998b; see also the discussion in Paper II).

The photometric data are presented in Table 3. Only the first page of this table is presented here for guidance regarding its form and content; the entire table is available in digital form on the Journal Electronic Edition, or from the authors. The table is organized as follows:

Col. 1: Galaxy name: if the galaxy is listed in the UGC catalog (Nilson 1973), the UGC number is given; if not, our internal coding number, from the private galaxy catalog of Giovanelli & Haynes referred to as the AGC, is given.

Cols. 2 and 3: the NGC or IC number, if available, and the CGCG field and ordinal number within that field, if the galaxy is listed in the CGCG (Zwicky *et al.* 1963-1968). For galaxies in the Coma cluster that are not listed in either one of those catalogs, the Dressler number (Dressler 1980) is given, if available.

Cols. 4 and 5: Right Ascension and Declination, in 1950.0 epoch.

Col. 6: morphological type code, in the RC3 (de Vaucouleurs *et al.* 1991) scheme (-5 for E, -2 for S0, 0 for S0a). Morphological types were derived from joint inspection of the blue plates of the Palomar Observatory Sky Survey (POSS), and of the CCD images.

Col. 7: Heliocentric radial velocity V_{hel} , in km s^{-1} .

Col. 8: Measured effective radius r_e , in arcsec.

Col. 9: Effective radius corrected for seeing effects r_e^c , in arcsec.

Col. 10: Uncertainty on the effective radius ϵ_r , in arcsec.

Col. 11: Corrected metric effective radius R_e , in kiloparsec, obtained assuming the galaxy is at the distance indicated by the cluster redshift, in the CMB reference frame.

Col. 12: Measured effective surface brightness μ_e , in I band mag arcsec $^{-2}$.

Col. 13: Corrected effective surface brightness μ_e^c , in I band mag arcsec $^{-2}$. μ_e^c is corrected for the smearing effects of seeing, for extinction within our Galaxy, for the cosmological k-correction term, and for $(1+z)^4$ cosmological dimming.

Col. 14: Uncertainty on the effective surface brightness ϵ_μ , in mag arcsec $^{-2}$.

Col. 15: Measured apparent I band magnitude m_I , obtained extrapolating the surface brightness profile to infinity.

Col. 16: Corrected apparent I band magnitude m_I^c , obtained from the value listed in col. 15, and corrected for extinction within our Galaxy, and for the cosmological k-correction term. No internal extinction correction was applied.

Col. 17: Uncertainty on the measured magnitude, ϵ_m .

Col. 18: Absolute magnitude M_I , obtained assuming that the galaxy is at the distance indicated by the cluster redshift, in the CMB reference frame.

Col. 19: Quality codes, as follows:

First digit = Photometric code: 1 = observation obtained in photometric conditions, uncertainty in the zero-point determination <0.03 mag; 2 = observation obtained in photometric conditions, uncertainty in the zero-point determination 0.03–0.05 mag; 6 = observation obtained in non photometric conditions, uncertainty in the zero-point determination <0.03 mag; 7 = observation obtained in non photometric conditions, uncertainty in the zero-point determination 0.03–0.06

mag.

Second digit = Image quality code: 0 = image is OK; 1 = small amount of contamination from nearby stars or other galaxies; 2 = significant amount of contamination from nearby stars or other galaxies; 3 = iterative subtraction of the contaminating object required; 4 = flat-fielding of sub-standard quality, due to scattered light from a nearby bright star.

Third digit = Profile code: 1 = $r^{1/4}$ fit good at all radii; 2 = $r^{1/4}$ fit acceptable at all radii; 3 = $r^{1/4}$ fit good only in the core of the galaxy; 4 = the $r^{1/4}$ fit does not provide an acceptable description of the observed photometric profile.

If an asterisk appears in this column, special comments on the object are included at the foot of the table.

A small subset of these data has already been published in Paper I, limited to galaxies in the Coma and A2634 clusters. Since then, a few galaxies have been reobserved, and the data combined to obtain new measurements of the relevant photometric parameters. The data presented here therefore supersede that previous report.

6. Internal and external comparison

Repeated observations of the same galaxies provide an opportunity to evaluate the internal consistency of the photometric measurements. In total, repeated observations of 71 galaxies are available. The results of the comparison of effective radius, effective surface brightness, and total magnitude are presented in Figure 2, and they are listed in Table 4. Also listed in this table are the mean statistical uncertainties associated with the derivation of those parameters, multiplied by a $\sqrt{2}$ factor to allow a direct comparison with the rms scatter in the observations (assuming equal error contributions to the rms scatter). Good agreement is found between the separate measurements, with only few isolated significant discrepancies. These are generally due to the uncertainties introduced by the contamination from other nearby objects in the field, as discussed in section 4.1. There is also good general agreement between the rms scatter derived from the comparison and the estimated statistical uncertainties.

The errors on r_e and μ_e are highly correlated. Figure 3a shows $\Delta \log R_e$ versus $\Delta \mu_e$ for the 71 pairs of observations. The linear correlation coefficient between $\Delta \log R_e$ and $\Delta \mu_e$ is $R = 0.962$. Because r_e and μ_e enter the FP via the combination $\log R_e - 0.32 \mu_e$, the correlated nature of the errors on these two quantities translates into a predicted uncertainty on their combination of 0.013, to be compared with the observed scatter of 0.020 (see Figure 3b). Note however that a reduction of R by 10% (the approximate level of uncertainty on its determination) would translate in an increased predicted uncertainty on $\log R_e - 0.32 \mu_e$ to 0.025.

No large dataset of I band photometric observations of early-type galaxies is available for an external comparison of our photometric measurements. An indirect test can be however obtained using comparisons with similar measurements presented by Lucey *et al.* (1991), Saglia, Bender &

Dressler (1993b), Jørgensen, Franx & Kjaergaard (1995) for galaxies in the Coma cluster. Since none of these studies used I band for the photometric observations, it is necessary to assume standard colors for E and S0 galaxies, before comparisons can be made. The V band magnitudes from Lucey *et al.* (1991), the B band magnitudes from Saglia *et al.* (1993b), and the Gunn r band magnitudes from Jørgensen *et al.* (1995) have been transformed to I band magnitudes using V–I, B–I, and r–I colors of 1.31, 2.25, and 1.07, respectively (Fukugita *et al.* 1995), without taking into consideration the small differences between the brightest and the faintest objects introduced by the color-magnitude relation of early-type galaxies. Because of this, the expected rms scatter in the true colors of E and S0 galaxies around these mean values is expected to be $\simeq 0.2$ mag.

There are 57 galaxies in common with the sample of Lucey *et al.* (1991), 47 with the sample of Saglia *et al.* (1993b), and 106 with the sample of Jørgensen *et al.* (1995). Table 5 and the panels of Figure 4, 5, and 6 show the results of the comparisons of effective radius, effective surface brightness, and the combination $\log R_e - 0.32 \mu_e$. In these figures, the difference (us - literature) is plotted vs. the average value we can derive for the various parameters using the two respective measurements. There are significant differences in the determinations of both the effective radius and the effective surface brightness. For example, differences as high as 2 mag in the determination of μ_e , after the color correction described above were applied, are present. However it is possible to show that the main source of these differences lies in the different procedures adopted to perform the $r^{1/4}$ fit, and in particular, in the choice of the portion of the photometric profile used for the fit. In this work, the fit was extended to an outer radius chosen on the basis of the shape of the photometric profile: for profiles that do not follow the $r^{1/4}$ relation at all radii, only the core of the galaxy was fitted. If the fit were to have been performed using all the points in the photometric profile, the fitted profile would have a different slope, and both the effective radius and the effective surface brightness would have different values. Another possible source of discrepancy is the choice of the inner radius from where to start the $r^{1/4}$ fit. In this work, this radius was chosen to be twice the radius of the seeing disk. A different choice would likewise lead to a fitted profile with a somewhat different slope, especially for objects with small effective radius.

To quantify this effect, we have reanalyzed the galaxies that show a discrepancy $|\Delta\mu_e| > 1$ mag in Figure 4 and 6 (a total of 20 galaxies). Considering these objects only, we observe differences between our measurements and those taken from the literature of $\Delta \log R_e = 0.48 \pm 0.33$ and $\Delta\mu_e = 1.46 \pm 1.13$, respectively. We have then redetermined r_e and μ_e for all these galaxies without using the $r^{1/4}$ fit. To do so, we measured on the observed photometric profile the radius that includes half of the total galaxy flux (as derived from the total magnitude, which, in turn, was obtained extrapolating to infinity the outer parts of the photometric profile), and the corresponding average surface brightness. The values of r_e and μ_e determined using this procedure are in much better agreement with those published by Saglia *et al.* (1993b), and by Jørgensen *et al.* (1995): the new differences between our modified measurements and those taken from the literature are $\Delta \log R_e = 0.063 \pm 0.084$ and $\Delta\mu_e = 0.086 \pm 0.29$, respectively. To help visualize the extent of this problem with different fitting techniques, the panels of Figure 4, 5, and 6 show with

filled symbols the galaxies whose photometric profiles follow the $r^{1/4}$ relation at all radii, and with unfilled symbols those galaxies that do not. It is clear that the largest discrepancies are mostly associated with the latter objects.

As we have shown, the observed differences in $\log R_e$ and μ_e originate from differences in the $r^{1/4}$ fit. As a result there is a correlation between $\Delta \log R_e$ and $\Delta \mu_e$ that is completely analogous to the correlation shown by the errors in $\log R_e$ and μ_e . Therefore, despite the large differences in the two separate values of $\log R_e$ and μ_e , we observe an excellent agreement in the comparison of the combinations $\log R_e - 0.32 \mu_e$. It is therefore possible to say that these external comparisons show good overall agreement, and establish the good quality of the photometric measurements presented in this work, as well as those presented in the comparison samples.

We would like to thank Gary Wegner, Shoko Sakai, and Bob Barr for their help with the MDM observations, and Bill Schoening for his technical support during the various observing runs at Kitt Peak. This research has made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration. This research is part of the Ph.D. Thesis of MS, and is supported by the NSF grants AST94–20505 and AST96–17069 to RG and AST95–28860 to MPH.

REFERENCES

- Bothun, G.D., Geller, M.J., Beers, T.C., & Huchra, J.P. 1983, *ApJ*, 268, 47
- Burstein, D., & Heiles, C. 1978, *ApJ*, 225, 40
- David, L.P., Slyz, A., Jones, C., Forman, W., Vrtillek, S.D., & Arnaud, K.A. 1993, *ApJ*, 412, 479
- de Vaucouleurs, G. 1948, *Ann. Astrophys.*, 11, 247
- de Vaucouleurs, G., de Vaucouleurs, A., Corwin, H.G., Buta, R., Paturel, G., & Fouqué, P. 1991, “Third Reference Catalogue of Bright Galaxies” (New York: Springer) (RC3)
- Djorgovski, S.G., & Davis, M. 1987, *ApJ*, 313, 59
- Djorgovski, S.G., De Carvalho, R.R., & Han, M.S. 1988, in “The Extragalactic Distance Scale”, ed. S. van den Bergh & J. Pritchet, p. 329
- Dressler, A. 1980, *ApJS*, 42, 565
- Dressler, A., Lynden-Bell, D., Burstein, D., Davies, R.L., Faber, S.M., Terlevich, R.J., & Wegner, G. 1987, *ApJ*, 313, 42
- Edge, A.C., & Stewart, G.C. 1991, *MNRAS*, 252, 428
- Forman, W., & Jones, C. 1982, *ARA&A*, 20, 547
- Fukugita, M., Shimasaku, K., & Ichikawa, T. 1995, *PASP*, 107, 945

- Gavazzi, G., Randone, I., & Branchini, E. 1995, *ApJ*, 438, 590
- Giovanelli, R., & Haynes, M.P. 1985, *AJ*, 90, 2445
- Giovanelli, R., Haynes, M.P., Herter, T., Vogt, N.P., da Costa, L.N., Freudling, W., Salzer, J.J., & Wegner, G. 1997a, *AJ*, 113, 22 (G97a)
- Giovanelli, R., Haynes, M.P., Herter, T., Vogt, N.P., da Costa, L.N., Freudling, W., Salzer, J.J., & Wegner, G. 1997b, *AJ*, 113, 53 (G97b)
- Guzmán, R., Lucey, J.R., Carter, D., & Terlevich, R.J. 1992, *MNRAS*, 257, 187
- Guzmán, R., Lucey, J.R., Bower, R.G., 1993, *MNRAS*, 265, 731
- Haynes, M.P., & Giovanelli, R., 1984, *AJ*, 89, 758
- Jørgensen, I., Franx, M. & Kjaergaard, P. 1995, *MNRAS*, 273, 1097
- Jørgensen, I., Franx, M., & Kjaergaard, P. 1996, *MNRAS*, 280, 167
- Kaiser, N. 1987, *MNRAS*, 227, 1
- Kogut, A., et al. 1993, *ApJ*, 419, 1
- Landolt, A.U. 1992, *AJ*, 104, 340
- Lucey, J.R., Guzmán, R., Carter, D., & Terlevich, R.J. 1991, *MNRAS*, 253, 584
- Magri, C., Haynes, M.P., Forman, W., Jones, C., & Giovanelli, R. 1988, *ApJ*, 333, 136
- Nilson, P. 1973, “Uppsala General Catalogue of Galaxies” (Uppsala Astron. Obs. Ann., Vol. 6) (UGC)
- Pahre, M.A., & Djorgovski, S.G., 1997, in “The Nature of Elliptical Galaxies, Proceedings of the Second Stromlo Symposium”, ASP Conf. Series 116, eds. M. Arnaboldi, G. S. Da Costa, and P. Saha, p. 154
- Pinkney, J., Rhee, G., Burns, J.O., Hill, J.M., Oegerle, W.M., Batuski, D., & Hintzen, P. 1993, *ApJ*, 416, 36
- Regös, E., & Geller, M.J. 1989, *AJ*, 98, 755
- Renzini, A., & Ciotti, L., 1993, *ApJ*, 416, L49
- Saglia, R., Bertschinger, E., Baggle, G., Burstein, D., Colless, M., Davies, R.L., McMahan, R.K., & Wegner, G. 1993a, *MNRAS*, 264, 961
- Saglia, R.P., Bender, R., & Dressler, A. 1993b, *A&A*, 279, 75
- Sakai, S., Giovanelli, R., & Wegner, G. 1994, *AJ*, 108, 33
- Sarazin, C.L., 1988, “X-ray emission from clusters of galaxies” (Cambridge: Cambridge University Press)
- Schindler, S., & Prieto, M.A., 1997, *A&A*, 327, 37
- Schlegel, D.J., Finkbeiner, D.P., & Davis, M., 1998, *ApJ*, 500, 525

- Schroeder, A., 1996, PhD Thesis, Univ. of Basel
- Scodeggio, M., Solanes, J.M., Giovanelli, R., & Haynes, M.P. 1995, *ApJ*, 444, 41
- Scodeggio, M., Giovanelli, R., & Haynes, M.P. 1997a, *AJ*, 113, 101 (Paper I)
- Scodeggio, M., Giovanelli, R., & Haynes, M.P. 1997b, *AJ*, 113, 2087 (Paper II)
- Scodeggio, M., Giovanelli, R., & Haynes, M.P. 1998a, submitted to *AJ*
- Scodeggio, M., Giovanelli, R., & Haynes, M.P. 1998b, in preparation
- Tully, R.B., & Fisher, J.R. 1977, *A&A*, 54, 661
- van Dokkum, P.G., & Franx, M. 1996, *MNRAS*, 281, 985
- van Haarlem, M.P. 1992, PhD Thesis, Univ. of Leiden
- Zabludoff, A.I., Huchra, J.P., & Geller, M.J. 1990, *ApJS*, 74, 1
- Zwicky, F., Herzog, E., Wild, P., Karpowicz, M., & Kowal, C. 1961–1968, “Catalogue of Galaxies and Clusters of Galaxies” (Pasadena: California Institute of Technology)

Table 2: Imaging observing runs

	Run 1	Run 2, 4–8	Run 3
Telescope	MDM 1.3m	KPNO 0.9m	MDM 1.3m
CCD	Loral2048 (Wilbur)	T2KA	Tek1024 (Charlotte)
Pixels	1024 x 1024	2048 x 2048	1024 x 1024
Read-out noise	6.0 e [−]	4.0 e [−]	4.8 e [−]
Gain	1.9 e [−] /DU	3.2 e [−] /DU	3.45 e [−] /DU
Pixel scale	0.63 ″/pixel	0.68 ″/pixel	0.51 ″/pixel
Field of view	11 x 11 arcmin	23 x 23 arcmin	9 x 9 arcmin

	Run 1	Run 2	Run 3	Run 4
Dates	Sep. 23-25 1992 ^a	Aug. 18-20 1993 ^b	Sep. 8-9 1993 ^a	Apr. 14-19 1994
Photometric nights	3	0	2	3
Marginal nights	0	3	0	1
Nr. of galaxies	10	78	6	322

	Run 5	Run 6	Run 7	Run 8
Dates	Aug. 31 - Sep. 6 1994	Oct. 8 1994 ^a	Mar. 6 1995 ^a	Sep. 24 - Oct. 1 1995
Photometric nights	3	1	0	2
Marginal nights	1	0	1	2
Nr. of galaxies	223	16	5	47

^aObserving run mainly dedicated to a different project; only a small portion of telescope time dedicated to this project.

^bService observing by KPNO personnel.

Table 4. Imaging data quality: internal comparison

Parameter	N_{gal}	mean difference	rms scatter	mean uncertainty
$\log r_e$	71	-0.003	0.044	0.038
μ_e	71	-0.001	0.131	0.113
m_{tot}	71	0.071	0.087	0.084
$\log r_e - 0.32\mu_e$	71	0.001	0.020	0.013

Table 5. Imaging data quality: External Comparison

Parameter	N_{gal}	mean difference	rms scatter
Saglia et al. (B band)			
$\log r_e$	47	0.158	0.240
μ_e	47	0.365	0.813
$\log r_e - 0.32\mu_e$	47	0.034	0.043
Lucey et al. (V band)			
$\log r_e$	57	0.030	0.100
μ_e	57	-0.100	0.353
$\log r_e - 0.32\mu_e$	57	0.060	0.030
Jørgensen et al. (r band)			
$\log r_e$	106	0.173	0.248
μ_e	106	0.438	0.741
$\log r_e - 0.32\mu_e$	106	0.031	0.038




Fig. 1.— The distribution of effective seeing for the galaxy observations. The pixel size of 0.68 arcsec. limits this effective seeing to be no smaller than $\simeq 1.2$ arcsec.




Fig. 2.— Internal comparison of the global photometric parameters. Δm_I , $\Delta \log_{10} R_e$, and $\Delta \mu_e$ are the differences between the values of the total magnitude, effective radius, and effective surface brightness obtained from repeated observations of 71 galaxies. They are plotted as a function of the mean value of the relative parameter, obtained combining the different measurements.




Fig. 3.— (a) The correlation of the errors in $\log r_e$ and μ_e . The figure shows the differences measured from repeated observations. The linear correlation coefficient for this data-set is 0.962. (b) Internal comparison of the combination $\log r_e - 0.32 \mu_e$, which is the combination that enters in the FP relation. As in Fig.2, the difference derived from repeated observations is plotted against the mean value of the parameter.




Fig. 4.— External comparison of the global photometric parameters for galaxies in the Coma cluster with the B band data from Saglia et al. B magnitudes were converted to I magnitudes for the comparison, assuming a fixed B-I color (see text). Filled symbols identify the galaxies whose photometric profile follows the $r^{1/4}$ relation at all radii, while empty symbols identify those galaxies that deviate significantly from such a profile. The differences (us - Saglia) are plotted against the mean value of the different parameters, obtained combining the two measurements.




Fig. 5.— External comparison of the global photometric parameters for galaxies in the Coma cluster with the V band data from Lucey et al. V magnitudes were converted to I magnitudes for the comparison, assuming a fixed V-I color (see text). Filled symbols identify the galaxies whose photometric profile follows the $r^{1/4}$ relation at all radii, while empty symbols identify those galaxies that deviate significantly from such a profile. As in the previous figure, differences (us - Lucey) are plotted against the mean value of the different parameters.




Fig. 6.— External comparison of the global photometric parameters for galaxies in the Coma cluster with the r band data of Jørgensen et al. r magnitudes were converted to I magnitudes for the comparison, assuming a fixed r-I color (see text). Filled symbols identify the galaxies whose photometric profile follows the $r^{1/4}$ relation at all radii, while empty symbols identify those galaxies that deviate significantly from such a profile. As in the previous figure, differences (us - Jørgensen) are plotted against the mean value of the different parameters.

TABLE 1. Clusters and Groups properties

Name	R.A. (1950)	Dec (1950)	V_{hel} — km s^{-1} —	V_{CMB}	σ_v	Richness	pop. fractions E:S0:S	L_X^a ergs s^{-1}	T_g keV	HI def %	Ref.
N383	01 04 39.4	+32 08 46	5161	4865	553	...	14:34:52 ^b	$<2.0 \cdot 10^{42}$	...	10 ^b	1
N507	01 20 51.1	+32 59 40	5091	4808	643	...	14:34:52 ^b	$<2.0 \cdot 10^{42}$	...	10 ^b	1
A262	01 49 49.9	+35 54 21	4896	4642	462	0	21:32:47	$2.63 \cdot 10^{43}$	2.4	48	2
Cancer ^c	08 17 40.7	+21 13 36	4705	4939	317	...	11:18:71	$<2.0 \cdot 10^{42}$	...	21	3
A1367	11 41 57.0	+20 03 30	6408	6735	822	2	17:40:43	$7.01 \cdot 10^{43}$	3.7	42	2
Coma	12 57 11.1	+28 13 48	6917	7185	1010	2	30:56:14	$6.02 \cdot 10^{44}$	8.3	77	2
Pegasus	23 17 42.6	+07 55 57	3888	3519	393	...	12:29:59	$<6.3 \cdot 10^{42}$	...	18	4
A2634	23 35 59.1	+26 45 14	9240	8895	661	1	17:46:37	$5.22 \cdot 10^{43}$	3.4	60	5

^a L_X measured in the 2–10 keV band.

^bComputed considering NGC 383 and NGC 507 together.

^cGroup A of Bothun et al. (1983).

References for Table 1.

(1) Sakai et al. 1994; (2) Zabludoff et al. 1990; (3) Bothun et al. 1983; (4) Giovanelli et al. 1997a; (5) Scodeggio et al. 1995

TABLE 3. Measured photometric parameters

Name	NGC	CGCG	R.A. (1950)	Dec (1950)	T	V_{hel} (km/s)	r_e ($''$)	r_e^c ($''$)	ϵ_r ($''$)	R_e (kpc)	μ_e – (mag/arcsec ²) –	μ_e^c (13)	ϵ_μ (14)	m_I (mag)	m_I^c (mag)	ϵ_m (mag)	M_I (mag)
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)
A110065	...	...	1 04 05.8	32 26 03	-5	5945	24.81	24.69	1.51	5.82	21.95	21.76	0.07	14.15	14.03	0.05	-19.40
A111066	...	...	1 04 12.6	32 02 30	-5	5581	2.44	1.87	0.11	0.44	17.09	16.66	0.07	13.46	13.36	0.04	-20.08
U680	N374	501-080	1 04 19.6	32 31 46	0	5031	22.36	22.23	0.54	5.24	19.89	19.70	0.04	11.45	11.34	0.05	-22.10
A110036	N375	...	1 04 20.2	32 04 52	-5	6011	2.28	1.70	0.13	0.40	16.60	16.13	0.09	13.34	13.23	0.04	-20.20
U683	N379	501-082	1 04 30.0	32 15 16	-2	5556	30.65	30.62	0.89	7.22	19.67	19.49	0.04	11.33	11.22	0.05	-22.21
U682	N380	501-081	1 04 32.3	32 13 01	-5	4414	7.94	7.69	0.12	1.81	17.69	17.49	0.03	10.93	10.83	0.05	-22.61
A111067	...	...	1 04 32.4	32 09 27	-2	...	9.82	9.61	0.70	2.27	20.85	20.66	0.15	14.86	14.76	0.10	-18.68
U688	N382	501-086	1 04 38.7	32 08 13	-5	5228	4.07	3.70	0.12	0.87	17.31	17.05	0.15	12.50	12.40	0.10	-21.04
U689	N383	501-087	1 04 39.4	32 08 46	-2	5071	28.05	27.99	0.31	6.60	19.39	19.21	0.03	9.59	9.49	0.06	-23.95
U686	N384	501-084	1 04 39.7	32 01 33	-2	4287	12.14	11.95	0.28	2.82	18.74	18.55	0.04	11.56	11.46	0.05	-21.98
U687	N385	501-085	1 04 41.9	32 03 15	-5	5024	10.67	10.47	0.20	2.47	18.74	18.55	0.03	11.43	11.33	0.05	-22.11
A110044	N386	501-088	1 04 45.5	32 05 43	-5	5540	11.96	11.77	0.49	2.78	19.68	19.49	0.15	13.20	13.09	0.10	-20.34
A111068	...	...	1 04 47.4	32 07 27	-5	4824	3.33	2.88	0.18	0.68	18.24	17.95	0.15	13.91	13.81	0.10	-19.63
A111069	...	...	1 05 00.5	32 25 34	-5	5434	7.67	7.37	0.44	1.74	19.72	19.50	0.08	13.76	13.65	0.05	-19.79
A110047	N388	501-090	1 05 01.4	32 02 36	-2	5497	5.70	5.41	0.21	1.28	18.40	18.18	0.06	12.95	12.84	0.05	-20.59
A111070	...	...	1 05 06.3	31 59 22	-5	...	14.99	14.85	1.21	3.50	21.73	21.54	0.10	15.06	14.96	0.05	-18.48
U700	N392	501-094	1 05 37.1	32 52 00	-2	4672	13.91	13.71	0.21	3.23	18.77	18.58	0.03	11.27	11.17	0.05	-22.27
A100055	N394	501-095	1 05 39.7	32 52 53	-2	4310	17.39	17.24	1.28	4.07	19.49	19.31	0.15	12.70	12.60	0.10	-20.84
A110057	N397	501-096	1 05 44.8	32 50 34	-2	4934	3.86	3.39	0.15	0.80	17.95	17.67	0.06	13.18	13.08	0.05	-20.36
U712	N399	501-101	1 06 13.0	32 22 06	0	5167	13.75	13.66	0.44	3.22	19.70	19.51	0.05	12.15	12.04	0.06	-21.39
U715	N403	501-104	1 06 28.2	32 29 05	0	4977	91.86	91.86	1.75	21.67	21.15	20.97	0.03	11.25	11.14	0.05	-22.29
U730	N407	501-115	1 07 49.9	32 51 32	-5	5573	40.06	40.06	1.36	9.45	20.32	20.14	0.04	11.80	10.58	0.05	-22.86
A111071	...	...	1 08 04.0	32 50 52	-5	...	10.18	10.06	0.36	2.37	20.59	20.40	0.05	13.75	13.64	0.06	-19.79
A111072	...	...	1 08 05.7	31 50 13	-2	...	5.62	5.45	0.44	1.29	20.27	20.07	0.11	14.55	14.45	0.06	-18.98
U735	N410	501-118	1 08 11.6	32 53 10	-5	5294	31.15	31.15	0.31	7.35	19.48	19.30	0.03	10.11	10.00	0.05	-23.43
A110098	...	501-119	1 08 20.8	32 57 41	-5	4809	12.51	12.42	0.29	2.93	19.80	19.62	0.04	12.59	12.48	0.05	-20.96
A111073	...	...	1 08 39.0	33 19 32	-5	...	6.16	5.98	0.30	1.41	20.41	20.21	0.07	14.77	14.66	0.06	-18.77
A110111	I1636	501-125	1 08 48.0	33 06 00	-2	5632	5.10	4.90	0.17	1.16	18.38	18.17	0.05	13.02	12.91	0.05	-20.53
A111074	...	...	1 08 54.1	33 15 17	-2	...	6.51	6.34	0.36	1.50	19.46	19.26	0.07	13.56	13.45	0.05	-19.98
A111136	...	...	1 09 18.4	31 48 45	-2	4661	14.92	14.84	0.63	3.50	21.23	21.05	0.05	14.31	14.21	0.05	-19.22
U752	N420	501-127	1 09 23.6	31 51 28	-5	5025	16.33	16.26	0.23	3.84	19.29	19.15	0.03	11.25	11.15	0.05	-22.28
A110121	I1638	501-129	1 09 35.5	33 05 47	1	4649	10.42	10.30	0.24	2.43	19.42	19.23	0.04	12.37	12.26	0.05	-21.17

TABLE 3. (continued)

Name	NGC	CGCG	R.A. (1950)	Dec (1950)	T	V_{hel} (km/s)	r_e ($''$)	r_e^c ($''$)	ϵ_r ($''$)	R_e (kpc)	μ_e – (mag/arcsec ²)	μ_e^c – (mag/arcsec ²)	ϵ_μ –	m_I (mag)	m_I^c (mag)	ϵ_m (mag)	M_I (mag)
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)
U792	I1652	502-009	1 12 09.7	31 41 03	-2	5176	45.86	45.86	1.62	10.82	20.80	20.64	0.04	12.40	12.31	0.05	-21.13
A111075	...	...	1 12 30.1	31 49 19	-5	...	9.77	9.66	0.54	2.28	21.10	20.93	0.07	14.37	14.28	0.06	-19.16
U804	N447	502-013	1 12 49.8	32 48 10	0	5597	16.60	16.51	0.42	3.85	19.60	19.41	0.04	11.33	11.21	0.06	-22.20
A110205	N468	502-029	1 17 00.0	32 30 07	1	5172	39.50	39.50	1.72	9.21	21.67	21.50	0.05	13.23	13.13	0.05	-20.27
A110207	...	502-031	1 17 05.7	32 41 30	-5	4408	8.20	8.05	0.42	1.88	19.75	19.57	0.07	13.24	13.15	0.05	-20.26
A111076	...	...	1 17 22.2	32 31 19	-5	...	6.18	6.00	0.40	1.40	20.12	19.93	0.09	14.46	14.37	0.05	-19.04
U870	N472	502-034	1 17 40.4	32 26 55	-5	5271	7.11	6.95	0.19	1.62	18.19	18.00	0.04	12.07	11.97	0.05	-21.44
A110226	I1673	502-038	1 17 57.9	32 46 59	-5	5102	4.30	4.06	0.14	0.95	17.86	17.65	0.06	12.83	12.74	0.04	-20.67
A111077	...	...	1 18 09.3	33 28 29	-5	...	3.68	3.44	0.21	0.80	19.19	18.97	0.08	14.46	14.37	0.05	-19.04
U878	...	521-004	1 18 13.9	33 38 12	-2	4043	22.66	22.63	1.32	5.28	19.60	19.45	0.07	11.28	11.20	0.05	-22.21
A110232	...	502-043	1 18 16.9	33 07 02	-5	5186	8.81	8.65	0.28	2.02	19.38	19.21	0.05	12.80	12.71	0.06	-20.70
A110237	...	502-044	1 18 28.9	32 49 44	-5	5018	7.07	6.88	0.21	1.60	19.12	18.95	0.05	12.96	12.87	0.05	-20.54
A111078	...	...	1 18 29.7	32 53 49	-5	...	4.68	4.44	0.31	1.03	19.64	19.44	0.09	14.61	14.52	0.04	-18.89
A111079	...	...	1 18 53.0	32 48 25	-2	...	2.32	1.91	0.17	0.45	18.76	18.44	0.11	14.94	14.85	0.05	-18.56
A110246	I1679	502-048	1 18 55.5	33 14 07	0	5300	28.12	28.12	1.38	6.55	21.73	21.57	0.06	13.56	13.47	0.06	-19.94
A110248	I1680	502-049	1 18 58.0	33 01 27	0	4377	3.24	2.91	0.12	0.68	17.46	17.21	0.06	12.96	12.88	0.05	-20.53
U906	N483	502-050	1 19 07.0	33 15 41	0	5997	6.08	5.87	0.12	1.37	17.83	17.64	0.04	11.97	11.88	0.06	-21.53
A111080	...	...	1 19 12.1	33 14 54	-2	...	19.57	19.49	1.00	4.54	21.38	21.22	0.15	14.14	14.05	0.10	-19.36
A111081	...	...	1 19 22.9	33 11 17	-2	...	10.92	10.78	0.53	2.51	20.73	20.56	0.06	14.51	14.42	0.05	-18.99
A110268	...	521-011	1 19 59.4	33 42 27	0	5381	61.65	61.65	3.59	14.37	21.86	21.69	0.06	13.53	13.43	0.05	-19.98
A110272	...	...	1 20 02.7	33 15 23	-2	4132	1.85	1.43	0.10	0.33	16.99	16.58	0.09	14.42	14.32	0.06	-19.09
A111082	...	...	1 20 04.2	33 09 09	-2	...	59.26	59.26	2.18	13.81	22.43	22.27	0.04	13.48	13.38	0.05	-20.03
U920	N495	502-058	1 20 07.1	33 12 40	1	4114	13.20	13.11	0.34	3.06	19.19	19.01	0.04	11.86	11.76	0.06	-21.65
U926	N499	502-059	1 20 22.8	33 12 01	-5	4375	23.23	23.21	0.25	5.41	18.92	18.75	0.03	10.28	10.18	0.06	-23.23
A110284	N501	502-062	1 20 33.3	33 10 20	-5	4887	5.97	5.76	0.22	1.34	18.75	18.56	0.06	13.06	12.96	0.06	-20.45
A111084	...	...	1 20 33.6	33 03 16	-2	...	11.27	11.12	0.50	2.59	20.95	20.77	0.06	14.52	14.42	0.05	-18.99
A111085	...	...	1 20 34.4	33 07 15	-2	...	16.50	16.41	1.12	3.82	21.05	20.88	0.08	14.77	14.67	0.06	-18.74
A111086	...	...	1 20 37.9	33 05 26	-2	...	2.72	2.32	0.20	0.54	18.62	18.33	0.11	15.73	15.63	0.05	-17.78
U935	N504	502-064	1 20 39.1	32 56 38	-2	4288	9.29	9.12	0.20	2.13	18.41	18.22	0.04	12.06	11.96	0.05	-21.45
A111087	...	...	1 20 39.2	32 49 21	-5	...	3.48	3.17	0.17	0.74	18.24	17.99	0.08	13.91	13.81	0.05	-19.60
A110285	N503	502-065	1 20 39.3	33 04 16	-2	5978	6.76	6.55	0.23	1.53	19.01	18.81	0.05	12.11	12.01	0.06	-21.40
U938	N507	502-067	1 20 51.1	32 59 40	-5	4915	16.90	16.80	0.22	3.92	18.62	18.45	0.03	9.86	9.76	0.06	-23.66

TABLE 3. (continued)

Name	NGC	CGCG	R.A. (1950)	Dec (1950)	T	V_{hel} (km/s)	r_e ($''$)	r_e^c ($''$)	ϵ_r ($''$)	R_e (kpc)	μ_e – (mag/arcsec ²) –	μ_e^c (13)	ϵ_μ (14)	m_I (mag)	m_I^c (mag)	ϵ_m (mag)	M_I (mag)
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)
U939	N508	502-068	1 20 51.6	33 01 09	-5	5519	13.11	12.99	0.24	3.03	19.30	19.12	0.15	11.72	11.62	0.10	-21.79
A110290	...	...	1 20 52.5	32 56 32	-5	5158	1.23	0.79	0.06	0.18	15.83	15.20	0.08	13.97	13.87	0.04	-19.54
A111088	...	...	1 20 54.1	33 09 21	-2	...	29.04	29.03	1.51	6.77	22.26	22.09	0.06	14.19	14.09	0.06	-19.32
A110833	I1689	502-070	1 20 58.9	32 47 43	-5	4567	11.41	11.29	0.54	2.63	19.18	19.01	0.06	12.38	12.28	0.04	-21.13
A110293	I1690	502-071	1 21 00.5	32 53 45	-2	4537	17.52	17.42	0.88	4.06	20.06	19.89	0.06	13.14	13.04	0.05	-20.37
A111089	...	...	1 21 01.0	33 19 28	-5	...	7.15	6.99	0.24	1.63	19.54	19.34	0.05	13.48	13.38	0.07	-20.03
A110296	...	502-072	1 21 09.3	33 03 10	-2	4995	7.59	7.40	0.30	1.72	19.12	18.93	0.06	12.88	12.78	0.05	-20.63
A110302	...	502-074	1 21 18.0	32 31 00	-2	6024	20.79	20.73	1.41	4.83	21.31	21.14	0.08	14.00	13.89	0.04	-19.52
A111090	...	...	1 21 19.3	32 41 59	-2	...	7.53	7.36	0.45	1.71	20.80	20.62	0.08	14.90	14.80	0.05	-18.61
A111091	...	...	1 21 31.1	32 42 56	-5	...	2.07	1.63	0.21	0.38	18.86	18.48	0.16	15.49	15.39	0.04	-18.02
A111092	...	...	1 21 36.5	33 08 49	0	...	17.65	17.56	1.21	4.09	21.32	21.14	0.08	13.99	13.88	0.05	-19.53
A111093	...	...	1 21 38.2	33 04 27	-2	...	15.54	15.44	0.98	3.60	21.17	21.00	0.08	14.98	14.87	0.05	-18.54
U956	N515	502-077	1 21 48.8	33 12 45	-2	4152	11.11	11.00	0.24	2.56	18.71	18.53	0.04	11.75	11.65	0.06	-21.76
A110305	I1692	502-076	1 21 50.3	32 58 33	-2	5525	25.10	25.08	0.90	5.85	21.22	21.04	0.05	13.61	13.50	0.05	-19.91
U960	N517	502-079	1 21 54.2	33 10 08	-5	4168	13.97	13.88	0.28	3.23	18.64	18.46	0.04	11.49	11.39	0.06	-22.02
A111094	...	...	1 21 58.0	33 05 03	-2	...	13.36	13.26	0.96	3.09	21.81	21.62	0.09	15.13	15.03	0.06	-18.38
A111095	...	...	1 22 41.6	33 21 20	-2	...	47.15	47.15	2.02	10.99	21.91	21.73	0.05	13.45	13.35	0.06	-20.06
A111096	...	...	1 22 44.4	32 59 39	-5	...	4.21	3.97	0.24	0.93	19.32	19.09	0.08	14.59	14.49	0.05	-18.92
U988	N528	502-083	1 22 44.6	33 24 45	-2	4879	27.44	27.44	0.57	6.40	19.80	19.62	0.04	11.55	11.45	0.06	-21.96
A111097	...	...	1 22 46.0	34 28 36	-5	...	5.93	5.74	0.45	1.34	19.30	19.11	0.10	13.94	13.85	0.05	-19.56
U995	N529	521-023	1 22 50.0	34 27 14	-5	4862	13.64	13.55	0.18	3.16	18.53	18.37	0.03	10.96	10.87	0.05	-22.54
A111098	...	...	1 22 53.4	32 49 41	-5	...	7.60	7.43	0.37	1.73	20.06	19.86	0.07	14.05	13.95	0.05	-19.46
A111099	...	...	1 22 58.4	33 05 09	0	...	35.26	35.26	1.44	8.22	21.44	21.27	0.05	13.49	13.39	0.05	-20.02
A111100	...	...	1 23 05.9	33 10 39	-2	...	36.60	36.60	4.73	8.53	21.62	21.45	0.12	14.49	14.39	0.05	-19.02
A110325	...	502-084	1 23 22.9	33 08 44	-2	5117	17.36	17.29	0.88	4.03	20.20	20.04	0.07	12.83	12.79	0.05	-20.62
U1012	N531	521-024	1 23 28.4	34 29 40	1	4660	16.59	16.50	0.81	3.85	20.00	19.84	0.06	12.44	12.35	0.05	-21.06
A110327	N542	521-026	1 23 40.4	34 24 56	-2	4674	366.8	366.88	22.23	85.52	23.69	23.53	0.04	13.64	13.55	0.05	-19.86
A111101	...	...	1 45 45.1	35 01 31	-5	...	3.03	2.75	0.23	0.61	18.82	18.59	0.12	14.85	14.77	0.05	-18.53
A110536	...	522-008	1 46 07.2	34 59 09	0	4423	14.70	14.61	0.76	3.24	20.42	20.27	0.07	13.80	13.72	0.05	-19.58
U1269	...	522-009	1 46 11.0	34 44 05	-5	3855	23.21	23.19	0.54	5.14	20.84	20.69	0.04	12.35	12.27	0.06	-21.03
U1272	...	522-010	1 46 21.2	34 49 26	1	5005	6.51	6.34	0.13	1.41	18.11	17.94	0.04	12.28	12.20	0.05	-21.10
A110538	...	522-011	1 46 23.2	34 43 13	-2	3900	7.10	6.94	0.43	1.54	19.48	19.31	0.09	14.20	14.12	0.05	-19.18

TABLE 3. (continued)

Name	NGC	CGCG	R.A. (1950)	Dec (1950)	T	V_{hel} (km/s)	r_e ($''$)	r_e^c ($''$)	ϵ_r ($''$)	R_e (kpc)	μ_e – (mag/arcsec ²)	μ_e^c –	ϵ_μ –	m_I (mag)	m_I^c (mag)	ϵ_m (mag)	M_I (mag)
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)
A110540	...	522-012	1 46 25.3	35 59 27	1	4041	30.65	30.65	1.20	6.80	21.43	21.25	0.05	13.65	13.54	0.05	-19.77
U1277	...	522-014	1 46 30.6	35 12 16	-2	4238	26.54	26.54	1.02	5.89	20.11	19.96	0.05	11.83	11.75	0.09	-21.55
A111102	...	...	1 46 43.8	35 54 28	-5	...	5.46	5.28	0.39	1.17	20.19	19.97	0.10	14.69	14.57	0.05	-18.73
A111103	...	...	1 46 44.5	35 59 28	-5	...	6.03	5.87	0.45	1.30	20.33	20.12	0.10	14.72	14.60	0.06	-18.70
U1283	N679	522-015	1 46 48.3	35 32 15	-2	5054	10.58	10.44	0.12	2.32	18.19	18.00	0.03	11.05	10.94	0.04	-22.36
A111104	...	...	1 47 18.8	35 58 53	-2	5175	7.43	7.29	0.33	1.62	19.60	19.39	0.06	13.69	13.57	0.05	-19.73
A111105	...	...	1 47 22.5	35 15 39	-5	...	5.75	5.56	0.45	1.23	19.90	19.70	0.11	14.35	14.25	0.05	-19.05
A111106	...	...	1 47 26.5	35 44 10	-2	...	29.25	29.25	2.26	6.49	22.52	22.34	0.08	14.33	14.22	0.08	-19.08
A111107	...	...	1 47 28.1	36 05 35	-2	4932	33.54	33.54	2.54	7.44	21.68	21.49	0.08	14.26	14.14	0.06	-19.16
U1298	N687	522-017	1 47 37.7	36 07 25	-2	5147	16.12	16.02	0.19	3.55	18.98	18.79	0.03	10.97	10.85	0.05	-22.45
A111108	...	...	1 47 45.5	36 05 08	-5	...	5.75	5.54	0.36	1.23	19.91	19.69	0.09	14.35	14.23	0.05	-19.07
A111109	...	...	1 47 47.3	35 30 55	-2	...	34.71	34.71	1.56	7.70	22.52	22.34	0.05	14.35	14.24	0.05	-19.06
U1307	I1732	522-021	1 47 51.8	35 41 07	-5	4889	26.53	26.52	0.77	5.88	19.86	19.68	0.04	11.42	11.31	0.05	-21.99
U1308	...	522-022	1 47 55.3	36 01 45	-2	5312	39.01	39.01	0.51	8.65	20.64	20.45	0.15	10.11	9.99	0.10	-23.31
A111110	...	...	1 48 00.6	35 53 52	-5	...	3.81	3.55	0.29	0.79	19.24	18.99	0.11	14.74	14.62	0.05	-18.68
A111111	...	...	1 48 12.8	36 00 43	-5	...	4.32	4.08	0.27	0.90	19.53	19.30	0.09	14.57	14.45	0.05	-18.85
A110851	...	522-023	1 48 23.6	34 37 29	0	5123	26.91	26.91	0.94	5.97	21.27	21.12	0.05	13.06	12.97	0.05	-20.33
A111112	...	...	1 48 30.4	35 41 16	-5	...	2.92	2.59	0.20	0.57	18.97	18.69	0.10	14.80	14.69	0.04	-18.61
A111113	...	...	1 48 36.5	35 37 44	-2	...	3.41	3.12	0.17	0.69	18.29	18.03	0.08	14.27	14.16	0.04	-19.14
A111114	...	...	1 48 41.5	36 20 37	-2	...	18.49	18.40	2.47	4.08	21.59	21.40	0.15	15.48	15.35	0.05	-17.95
A111115	...	...	1 48 52.7	34 48 11	-5	...	5.63	5.43	0.32	1.20	20.34	20.15	0.08	14.71	14.62	0.06	-18.68
A111116	...	...	1 48 56.4	35 25 00	-2	...	31.03	31.02	1.74	6.88	21.68	21.54	0.06	13.27	13.16	0.05	-20.14
A111117	...	...	1 49 02.0	34 45 55	-2	...	6.36	6.18	0.86	1.37	20.61	20.43	0.18	15.54	15.45	0.06	-17.85
A111118	...	...	1 49 09.0	34 39 01	-5	...	4.76	4.55	0.48	1.01	20.34	20.16	0.14	15.07	14.98	0.05	-18.32
A110581	...	522-026	1 49 11.1	35 10 04	-2	4877	6.94	6.78	0.21	1.50	18.78	18.58	0.05	12.97	12.86	0.05	-20.44
U1336	N700	522-027	1 49 16.4	35 51 03	-2	4364	13.65	13.55	0.47	3.00	20.33	20.12	0.05	13.02	12.89	0.05	-20.41
A110586	...	522-030	1 49 20.6	35 47 24	-2	4595	8.94	8.78	0.31	1.95	19.43	19.21	0.05	13.21	13.08	0.05	-20.22
U1339	...	522-032	1 49 28.7	35 35 35	1	4063	26.69	26.68	0.70	5.92	20.58	20.41	0.04	12.38	12.28	0.04	-21.03
A111119	...	...	1 49 31.2	36 15 07	-2	...	30.63	30.64	1.72	6.79	22.00	21.80	0.06	14.85	14.72	0.05	-18.59
A110848	N704B	522-034S	1 49 36.1	35 52 09	-5	...	6.03	5.82	0.19	1.29	19.04	18.81	0.15	13.37	13.24	0.10	-20.07
A111120	...	...	1 49 40.1	36 24 35	-5	...	4.22	3.98	0.28	0.88	19.99	19.74	0.09	15.08	14.95	0.05	-18.36
U1343N	N704A	522-034N	1 49 41.2	35 52 45	-5	4618	8.86	8.70	0.33	1.93	18.82	18.60	0.15	12.43	12.30	0.10	-21.01
A111121	...	...	1 49 42.6	35 54 30	-5	...	5.32	5.10	0.38	1.13	20.45	20.22	0.15	15.04	14.91	0.10	-18.40

TABLE 3. (continued)

Name	NGC	CGCG	R.A. (1950)	Dec (1950)	T	V_{hel} (km/s)	r_e ($''$)	r_e^c ($''$)	ϵ_r ($''$)	R_e (kpc)	μ_e – (mag/arcsec ²)	μ_e^c – (mag/arcsec ²)	ϵ_μ ($''$)	m_I (mag)	m_I^c (mag)	ϵ_m (mag)	M_I (mag)
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)
U1346	N703	522-037	1 49 43.2	35 55 28	-5	5175	14.69	14.59	0.50	3.24	19.20	19.00	0.15	11.75	11.61	0.10	-21.69
A111122	...	...	1 49 43.6	35 53 42	-5	...	2.15	1.72	0.17	0.38	17.49	17.10	0.15	14.49	14.36	0.10	-18.95
U1345	N705	522-036	1 49 45.4	35 53 51	0	4526	27.01	27.00	1.14	5.99	19.97	19.77	0.15	12.44	12.31	0.10	-20.99
U1348	N708	522-039	1 49 49.9	35 54 22	-5	4898	61.62	61.62	1.39	13.66	21.11	20.91	0.04	9.12	8.99	0.12	-24.32
A110590	N709	522-040	1 49 54.1	35 58 35	-5	3338	9.52	9.36	0.44	2.08	18.91	18.70	0.07	12.61	12.48	0.11	-20.82
A111123	...	...	1 49 54.3	35 54 20	-5	...	5.53	5.31	0.34	1.18	20.02	19.79	0.15	14.70	14.57	0.10	-18.74
U1343S	N704B	522-034S	1 49 00.0	35 52 00	-5	...	4.40	4.16	0.20	0.92	18.18	17.94	0.15	13.17	13.04	0.10	-20.27
U1352	N712	522-043	1 50 11.9	36 34 32	-2	5286	47.72	47.72	0.80	10.58	20.56	20.37	0.03	11.40	11.28	0.04	-22.02
A111124	...	...	1 50 12.7	36 06 01	-2	...	10.34	10.19	0.40	2.26	20.07	19.86	0.06	13.88	13.75	0.05	-19.57
A111125	...	...	1 50 13.1	36 39 39	-2	...	19.64	19.56	1.43	4.34	21.93	21.74	0.08	15.29	15.17	0.04	-18.13
A110594	...	522-044	1 50 18.1	36 05 17	-2	4984	12.21	12.10	0.47	2.68	19.89	19.69	0.05	13.34	13.21	0.05	-20.10
A111126	...	...	1 50 25.0	36 06 53	-2	...	40.59	40.59	3.19	9.00	22.49	22.29	0.08	15.10	14.97	0.05	-18.34
U1353	...	522-046	1 50 25.8	36 42 35	-5	5055	6.98	6.80	0.15	1.51	18.24	18.03	0.04	12.15	12.03	0.04	-21.27
A110597	...	522-045	1 50 27.3	35 45 57	-5	5113	23.72	23.69	0.80	5.25	20.88	20.68	0.05	12.18	12.04	0.09	-21.26
U1358	N714	522-047	1 50 32.9	35 58 33	0	4470	30.70	30.71	0.80	6.81	19.80	19.60	0.04	11.68	11.55	0.05	-21.75
A111127	...	...	1 50 44.7	36 04 06	-5	...	4.36	4.11	0.26	0.91	19.46	19.22	0.09	14.50	14.37	0.05	-18.95
A110605	...	522-048	1 50 53.3	36 06 15	-5	4114	10.67	10.53	0.23	2.34	19.32	19.11	0.04	12.34	12.21	0.05	-21.10
U1363	N717	522-052	1 50 58.3	35 59 01	0	4968	41.05	41.05	2.45	9.10	20.84	20.64	0.07	12.61	12.48	0.05	-20.83
A110607	...	522-053	1 50 59.9	36 23 42	1	4430	32.66	32.66	1.54	7.24	21.29	21.10	0.06	13.34	13.22	0.05	-20.09
A110608	...	522-054	1 51 14.8	35 40 18	1	4293	98.99	98.99	4.58	21.95	22.69	22.52	0.04	13.62	13.51	0.04	-19.79
A111128	...	...	1 51 54.4	36 13 06	-5	...	6.58	6.39	0.29	1.42	20.21	20.00	0.06	14.19	14.06	0.05	-19.24
A110613	...	522-057	1 51 55.6	37 17 01	0	4615	17.19	17.13	1.50	3.80	20.57	20.43	0.07	13.04	12.96	0.05	-20.34
A111129	...	...	1 51 57.5	36 04 40	-5	...	5.18	4.94	0.47	1.10	19.93	19.69	0.13	14.87	14.73	0.05	-18.57
A111130	...	...	1 52 00.5	36 29 37	-5	...	9.72	9.59	0.50	2.13	20.85	20.66	0.07	14.01	13.90	0.05	-19.40
A111131	...	...	1 52 03.3	36 09 56	-2	...	58.20	58.20	2.51	12.91	22.82	22.63	0.05	13.77	13.64	0.05	-19.66
A111132	...	...	1 52 13.4	36 20 17	-2	...	7.45	7.28	0.36	1.61	19.71	19.51	0.07	14.38	14.27	0.05	-19.03
U1387	...	522-063	1 52 13.7	36 01 06	-5	4502	5.76	5.53	0.23	1.23	18.76	18.52	0.06	13.34	13.20	0.05	-20.10
A111133	...	...	1 52 22.5	37 03 44	-5	...	2.64	2.32	0.14	0.51	17.68	17.43	0.08	13.89	13.81	0.05	-19.49
U1390	...	522-066	1 52 24.3	36 03 42	-2	4380	17.43	17.33	0.61	3.84	20.01	19.80	0.05	13.42	13.28	0.05	-20.02
U1392	...	...	1 52 32.6	36 19 03	1	5278	53.42	53.42	1.80	11.84	22.08	21.90	0.04	13.25	13.14	0.05	-20.16
U1406	N732	522-076	1 53 30.1	36 33 32	0	5813	14.57	14.49	0.59	3.21	19.66	19.47	0.05	12.13	12.01	0.05	-21.29
U1415	...	522-080	1 53 46.5	36 08 17	0	4796	21.27	21.23	0.72	4.71	19.90	19.72	0.05	12.54	12.42	0.04	-20.88
A110683	...	522-085	1 54 39.3	37 12 43	-5	5487	24.06	24.06	0.91	5.34	21.29	21.14	0.04	13.28	13.20	0.05	-20.10

TABLE 3. (continued)

Name	NGC	CGCG	R.A. (1950)	Dec (1950)	T	V_{hel} (km/s)	r_e ($''$)	r_e^c ($''$)	ϵ_r ($''$)	R_e (kpc)	μ_e – (mag/arcsec ²) –	μ_e^c (13)	ϵ_μ (14)	m_I (mag)	m_I^c (mag)	ϵ_m (mag)	M_I (mag)
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)
U1434	...	522-084	1 54 41.5	36 00 41	0	4564	36.34	36.34	1.40	8.06	20.95	20.74	0.05	13.03	12.89	0.04	-20.41
U1440	N759	522-087	1 54 53.1	36 06 00	-5	4714	16.02	15.94	0.21	3.53	19.09	18.91	0.03	11.12	11.00	0.04	-22.30
A110690	...	522-091	1 55 06.9	37 20 06	-5	4669	4.97	4.78	0.13	1.06	18.12	17.94	0.05	12.81	12.73	0.05	-20.57
A111135	...	...	1 55 35.5	35 58 23	-2	...	16.11	16.03	2.09	3.55	20.32	20.12	0.15	14.03	13.90	0.04	-19.41
U1475	I179	522-101	1 57 12.3	37 46 43	-5	4209	11.28	11.19	0.16	2.48	18.51	18.34	0.03	11.31	11.21	0.05	-22.09
A180152	I2253	119-024	8 13 38.3	21 33 53	-2	4741	8.64	8.49	0.18	2.03	18.43	18.26	0.04	12.17	12.09	0.05	-21.38
A180170	N2553	119-031	8 14 40.0	21 03 32	-2	4721	13.89	13.80	0.32	3.30	19.35	19.18	0.04	11.91	11.82	0.06	-21.65
A180195	N2556	119-045	8 16 05.9	21 05 39	-5	4635	5.41	5.00	0.18	1.20	18.25	18.02	0.05	12.97	12.89	0.05	-20.58
U4330	N2557	119-048	8 16 15.2	21 35 34	-2	4854	19.26	19.18	0.37	4.59	19.49	19.34	0.03	11.81	11.73	0.05	-21.73
A180199	...	119-053	8 16 24.8	21 12 58	-2	4850	23.58	23.44	1.75	5.61	21.52	21.35	0.08	14.01	13.92	0.06	-19.55
A180200	...	...	8 16 30.0	21 10 00	-5	3738	48.67	48.68	3.68	11.66	22.80	22.64	0.08	14.22	14.14	0.07	-19.33
U4337	N2560	119-058	8 16 57.0	21 08 34	-2	4928	28.58	28.46	0.70	6.81	20.01	19.85	0.04	11.15	11.06	0.07	-22.40
U4345	N2562	119-063	8 17 28.6	21 17 27	-2	4963	38.89	38.89	1.28	9.31	19.94	19.78	0.04	11.48	11.39	0.05	-22.08
U4347	N2563	119-065	8 17 40.7	21 13 40	-5	4642	21.73	21.64	0.22	5.18	19.19	19.03	0.03	10.36	10.28	0.07	-23.19
A180220	...	...	8 17 54.0	21 13 00	-2	4552	1.94	1.42	0.10	0.34	16.74	16.26	0.08	13.66	13.58	0.06	-19.89
A180225	N2569	119-067	8 18 26.4	21 01 37	-5	5104	9.44	9.26	0.22	2.22	19.09	18.91	0.04	12.43	12.34	0.06	-21.13
A180226	...	119-069	8 18 31.4	21 17 25	-5	4952	3.54	3.20	0.15	0.77	18.34	18.10	0.06	13.89	13.80	0.05	-19.66
A180237	...	119-077	8 19 58.2	21 14 21	-2	4473	12.00	11.78	0.81	2.82	19.41	19.24	0.09	13.21	13.13	0.06	-20.34
U4384	I2341	119-081	8 20 46.3	21 35 49	-2	4846	12.26	12.13	0.19	2.90	19.00	18.84	0.03	12.05	11.97	0.06	-21.50
A210505	...	97-021	11 33 51.1	20 05 42	-2	6646	5.28	5.02	0.11	1.64	17.99	17.83	0.04	12.26	12.24	0.06	-21.91
A210512	...	97-023	11 34 15.2	20 16 54	-5	6328	4.66	4.38	0.20	1.43	18.46	18.29	0.07	13.30	13.28	0.05	-20.86
A211586	...	...	11 34 28.7	20 12 02	-5	...	1.86	1.34	0.09	0.44	16.88	16.42	0.08	14.18	14.16	0.05	-19.98
A211587	...	...	11 37 14.9	20 40 19	-2	...	42.53	42.53	2.71	13.89	22.29	22.17	0.07	14.23	14.21	0.06	-19.93
A211588	...	...	11 37 23.0	20 26 12	-2	...	1.73	1.21	0.10	0.40	16.48	15.98	0.09	14.02	14.00	0.05	-20.14
U6642	N3805	127-024	11 38 05.9	20 37 13	-2	6472	21.70	21.61	0.67	7.06	19.14	19.02	0.05	10.84	10.82	0.06	-23.32
A211589	...	...	11 38 19.2	20 36 06	-2	...	10.92	10.75	1.10	3.51	20.34	20.21	0.12	14.84	14.82	0.05	-19.32
A210603	...	97-055	11 38 46.1	20 14 27	-5	6554	8.24	8.04	0.27	2.63	18.99	18.85	0.05	12.57	12.55	0.05	-21.60
A211590	...	...	11 39 03.2	20 37 15	-5	...	5.15	4.91	0.37	1.60	19.74	19.59	0.10	14.33	14.31	0.06	-19.83
U6656	N3816	97-060	11 39 12.0	20 22 55	-2	5548	21.40	21.32	0.36	6.96	19.66	19.54	0.03	11.18	11.16	0.06	-22.98
A211591	...	...	11 39 16.7	20 31 35	-2	...	6.41	6.19	0.44	2.02	19.83	19.68	0.09	14.56	14.54	0.05	-19.60
A211592	...	...	11 39 23.6	20 31 57	-2	...	14.82	14.71	1.22	4.80	21.88	21.75	0.10	15.20	15.18	0.06	-18.96

TABLE 3. (continued)

Name	NGC	CGCG	R.A. (1950)	Dec (1950)	T	V_{hel} (km/s)	r_e ($''$)	r_e^c ($''$)	ϵ_r ($''$)	R_e (kpc)	μ_e – (mag/arcsec ²) –	μ_e^c (13)	ϵ_μ (14)	m_I (mag)	m_I^c (mag)	ϵ_m (mag)	M_I (mag)
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)
U6663	N3821	127-032	11 39 33.0	20 35 38	0	5600	8.89	8.72	0.21	2.85	18.55	18.41	0.04	11.85	11.83	0.05	-22.31
A211621	...	...	11 40 06.2	20 34 37	-5	...	4.57	4.31	0.23	1.41	19.12	18.95	0.07	14.25	14.23	0.06	-19.91
A211593	...	...	11 40 06.2	20 34 37	-5	...	4.60	4.35	0.24	1.42	19.13	18.97	0.08	14.21	14.19	0.06	-19.95
A211149	...	...	11 40 08.9	20 13 34	-2	...	18.19	18.08	1.32	5.90	21.61	21.49	0.08	14.51	14.49	0.06	-19.65
A211594	...	...	11 40 16.5	19 42 22	-5	...	13.08	12.96	0.66	4.23	20.85	20.72	0.06	13.56	13.54	0.07	-20.60
A210639	...	97-074	11 40 23.3	20 21 42	-2	6496	5.51	5.27	0.17	1.72	18.54	18.39	0.05	13.13	13.11	0.05	-21.03
A211595	...	...	11 40 24.7	20 29 09	-5	...	3.98	3.70	0.24	1.21	19.06	18.88	0.09	14.21	14.19	0.06	-19.95
A211596	...	...	11 40 30.7	20 12 59	-2	...	9.77	9.59	0.53	3.13	20.34	20.20	0.07	14.06	14.04	0.06	-20.10
A211597	...	...	11 40 39.4	19 42 14	-5	...	2.59	2.17	0.16	0.71	18.33	18.07	0.09	14.64	14.62	0.06	-19.52
U6683	...	97-078	11 40 41.2	20 01 32	-5	7526	19.66	19.56	0.89	6.39	20.22	20.09	0.06	12.15	12.12	0.08	-22.02
A210647	...	97-077	11 40 41.8	19 39 50	-5	6531	6.90	6.69	0.27	2.18	19.18	19.04	0.06	13.32	13.30	0.06	-20.85
A211598	...	...	11 40 44.1	19 42 01	-2	...	10.72	10.56	0.77	3.45	21.08	20.95	0.09	14.88	14.86	0.07	-19.28
A211599	...	...	11 40 59.8	19 26 08	-5	...	7.09	6.86	1.24	2.24	21.37	21.22	0.22	15.36	15.34	0.08	-18.80
A210661	...	97-085	11 41 01.0	19 53 02	-2	6564	5.02	4.78	0.22	1.56	18.97	18.81	0.07	13.51	13.49	0.06	-20.66
A210662	N3834	97-084	11 41 01.8	19 22 12	-2	6824	8.68	8.49	0.20	2.77	19.02	18.88	0.04	12.55	12.53	0.06	-21.62
A211600	...	...	11 41 03.6	19 51 25	-2	...	4.14	3.87	0.31	1.26	19.56	19.38	0.11	14.71	14.69	0.06	-19.45
A210666	...	127-040	11 41 07.4	20 32 47	-2	6906	15.69	15.59	0.52	5.09	19.93	19.81	0.05	13.08	13.06	0.05	-21.09
A210667	...	97-086	11 41 07.5	20 18 58	-2	6399	17.09	16.99	1.43	5.55	20.14	20.02	0.10	13.73	13.71	0.05	-20.43
A211601	...	...	11 41 10.0	19 40 12	-5	...	5.5	5.26	0.26	1.72	19.86	19.71	0.07	14.30	14.28	0.06	-19.86
A211602	...	...	11 41 18.0	20 01 00	-2	...	13.08	12.94	1.03	4.23	20.67	20.54	0.09	14.73	14.71	0.05	-19.44
U6701	N3837	97-089	11 41 21.0	20 10 21	-5	6248	6.66	6.42	0.10	2.09	17.99	17.85	0.03	11.94	11.92	0.05	-22.23
A211622	...	...	11 41 21.9	20 34 40	-5	...	8.62	8.44	0.36	2.76	19.62	19.48	0.06	13.44	13.42	0.06	-20.72
A211603	...	...	11 41 21.9	20 34 40	-5	...	8.24	8.06	0.38	2.63	19.49	19.35	0.07	13.40	13.38	0.06	-20.76
A211604	...	...	11 41 21.9	20 12 47	-2	...	7.71	7.50	0.54	2.45	20.26	20.12	0.15	14.89	14.87	0.10	-19.27
A210674	...	97-090B	11 41 21.9	20 14 00	-2	6040	5.17	4.91	0.12	1.60	18.26	18.10	0.15	13.44	13.42	0.10	-20.72
A210676	...	97-088	11 41 23.5	20 03 35	-2	5624	6.25	6.01	0.21	1.96	18.66	18.52	0.05	13.14	13.12	0.06	-21.02
A210679	N3841	97-096	11 41 24.0	20 14 27	-2	6363	7.47	7.25	0.16	2.37	18.72	18.58	0.15	12.20	12.18	0.10	-21.96
U6705	N3844	97-097	11 41 24.8	20 18 19	-5	6834	24.08	24.03	0.68	7.84	20.13	20.02	0.04	11.71	11.69	0.08	-22.46
U6704	N3842	97-095	11 41 26.4	20 13 40	-5	6237	26.67	26.63	0.28	8.70	19.70	19.58	0.03	10.63	10.61	0.06	-23.53
A210677	...	97-094	11 41 27.6	20 04 47	-5	8004	3.45	3.10	0.16	1.01	18.11	17.89	0.07	13.83	13.80	0.06	-20.34
A210683	N3845	97-100	11 41 29.9	20 16 35	-2	5640	14.42	14.31	0.56	4.67	19.98	19.85	0.05	12.53	12.51	0.06	-21.63
A211605	...	...	11 41 30.2	20 31 32	-2	...	6.84	6.63	0.41	2.16	19.54	19.39	0.08	14.35	14.33	0.06	-19.81
A210688	...	97-099	11 41 31.5	20 00 58	-2	7211	5.45	5.20	0.18	1.70	19.07	18.91	0.05	13.44	13.41	0.05	-20.73

TABLE 3. (continued)

Name	NGC	CGCG	R.A. (1950)	Dec (1950)	T	V_{hel} (km/s)	r_e ($''$)	r_e^c ($''$)	ϵ_r ($''$)	R_e (kpc)	μ_e – (mag/arcsec ²)	μ_e^c (mag/arcsec ²)	ϵ_μ ($''$)	m_I (mag)	m_I^c (mag)	ϵ_m (mag)	M_I (mag)
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)
A210694	...	97-101	11 41 43.7	20 07 29	0	6454	11.43	11.27	0.36	3.68	19.51	19.38	0.04	12.42	12.40	0.06	-21.74
A210695	...	97-105	11 41 44.7	20 06 24	0	5440	11.35	11.18	0.53	3.65	19.89	19.76	0.06	12.83	12.81	0.06	-21.33
A210691	N3851	97-106	11 41 44.8	20 15 33	-2	6469	9.65	9.47	0.25	3.09	19.20	19.06	0.04	12.95	12.93	0.05	-21.21
A210703	...	97-110	11 41 49.3	20 06 27	-5	4527	9.28	9.09	0.31	2.97	19.60	19.46	0.05	13.14	13.12	0.06	-21.02
A211606	...	...	11 41 49.6	20 23 07	-2	...	3.90	3.61	0.21	1.18	18.38	18.19	0.08	13.62	13.60	0.05	-20.54
A210699	...	97-109	11 41 52.8	20 00 46	-5	6823	1.59	1.08	0.10	0.35	16.01	15.47	0.10	13.65	13.62	0.05	-20.52
A210708	...	97-112	11 41 54.4	20 21 13	-2	6753	12.25	12.12	0.71	3.96	19.18	19.05	0.07	12.72	12.70	0.06	-21.45
A210717	...	97-113	11 42 11.1	20 02 05	-2	6419	5.65	5.40	0.26	1.76	18.89	18.74	0.07	14.01	13.99	0.05	-20.15
A210724	...	97-115	11 42 12.4	20 09 19	-5	7792	11.66	11.50	0.43	3.76	19.90	19.76	0.05	12.87	12.84	0.07	-21.30
A210719	...	...	11 42 12.7	19 51 32	-2	6556	16.19	16.09	0.86	5.25	20.90	20.78	0.07	13.95	13.93	0.06	-20.22
A211330	...	...	11 42 12.7	19 51 30	-5	6225	8.66	8.48	0.53	2.77	20.56	20.42	0.08	13.99	13.97	0.06	-20.17
A210725	N3857	97-117	11 42 14.3	19 48 33	1	6256	14.81	14.70	0.32	4.80	19.99	19.86	0.04	12.32	12.30	0.06	-21.84
A210727	...	97-125	11 42 19.0	20 03 33	-5	8248	7.64	7.43	0.36	2.43	19.52	19.37	0.07	13.28	13.25	0.06	-20.89
A210729	...	97-124	11 42 20.7	20 00 11	-2	...	4.70	4.42	0.17	1.44	18.52	18.35	0.06	13.23	13.21	0.05	-20.93
A211608	...	...	11 42 27.2	20 58 20	-2	...	2.81	2.42	0.25	0.79	18.86	18.61	0.13	15.29	15.24	0.05	-18.90
U6723	N3862	97-127	11 42 29.1	19 53 05	-2	6462	11.50	11.36	0.19	3.71	18.80	18.67	0.03	10.34	10.32	0.09	-23.82
A210739	I2955	97-128	11 42 29.9	19 53 10	-2	6345	6.25	6.03	0.19	1.97	18.40	18.25	0.15	13.02	13.00	0.10	-21.14
A211609	...	...	11 42 35.3	19 48 38	-5	...	4.04	3.75	0.22	1.22	18.64	18.46	0.08	14.04	14.02	0.05	-20.12
A210741	...	97-131	11 42 38.7	20 07 25	-2	7646	13.73	13.62	0.35	4.45	19.78	19.64	0.04	12.53	12.50	0.06	-21.65
A211610	...	...	11 42 50.0	19 34 32	-5	...	4.19	3.89	0.29	1.27	19.52	19.34	0.10	14.51	14.49	0.06	-19.65
A211611	...	...	11 42 50.1	20 05 44	-2	...	4.97	4.75	0.51	1.55	18.90	18.75	0.14	14.65	14.63	0.05	-19.51
A210749	...	127-048	11 42 51.8	21 05 40	-5	6935	5.79	5.56	0.16	1.82	18.45	18.27	0.05	12.77	12.72	0.05	-21.42
A211612	...	...	11 42 52.7	20 34 18	-2	...	7.63	7.44	0.38	2.43	19.34	19.20	0.07	14.26	14.24	0.05	-19.90
A210750	N3868	97-135	11 42 54.6	19 43 29	-2	6653	16.73	16.62	0.38	5.43	19.69	19.57	0.04	12.17	12.15	0.06	-22.00
U6731	N3867	97-134	11 42 55.4	19 41 14	0	5000	24.68	24.62	0.56	8.04	20.33	20.21	0.04	11.52	11.50	0.06	-22.64
A211613	...	...	11 42 60.0	19 41 39	-5	...	10.05	9.87	0.55	3.22	20.77	20.63	0.07	13.85	13.83	0.07	-20.31
A211614	...	...	11 43 05.8	20 02 52	-5	...	4.02	3.77	0.21	1.23	19.03	18.86	0.08	14.11	14.09	0.05	-20.05
U6735	N3873	97-137	11 43 10.2	20 03 08	-5	5438	13.54	13.44	0.20	4.39	19.11	18.98	0.03	11.50	11.48	0.06	-22.66
U6739	N3875	97-139	11 43 13.8	20 02 24	-2	7088	38.34	38.34	0.84	12.52	20.49	20.37	0.15	12.45	12.42	0.10	-21.72
A210759	...	97-140	11 43 23.4	19 48 24	0	5486	5.51	5.26	0.22	1.72	18.66	18.51	0.06	13.12	13.10	0.05	-21.04
A210763	...	97-141	11 43 27.5	19 43 03	-2	5165	14.48	14.36	0.51	4.69	20.58	20.45	0.05	13.74	13.72	0.06	-20.42
A210764	...	97-142	11 43 29.9	19 55 06	-2	6357	17.78	17.68	0.86	5.77	20.83	20.71	0.06	14.04	14.02	0.06	-20.12
A210765	...	97-143A	11 43 29.9	20 04 11	-5	...	6.03	5.82	0.24	1.90	18.68	18.53	0.06	13.20	13.18	0.06	-20.96

TABLE 3. (continued)

Name	NGC	CGCG	R.A. (1950)	Dec (1950)	T	V_{hel} (km/s)	r_e ($''$)	r_e^c ($''$)	ϵ_r ($''$)	R_e (kpc)	μ_e – (mag/arcsec ²) –	μ_e^c (13)	ϵ_μ (14)	m_I (mag)	m_I^c (mag)	ϵ_m (mag)	M_I (mag)
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)
U6760	N3886	97-147	11 44 29.9	20 06 56	-5	5767	12.98	12.86	0.16	4.20	18.71	18.58	0.03	11.51	11.49	0.05	-22.65
A210795	...	97-155	11 45 28.0	20 17 10	-5	7280	6.71	6.50	0.20	2.12	18.43	18.27	0.05	12.62	12.59	0.05	-21.55
U6810	N3919	97-161	11 48 06.8	20 17 35	-5	6244	8.50	8.31	0.14	2.71	18.43	18.23	0.03	11.84	11.75	0.05	-22.39
A210831	...	97-162	11 48 16.0	20 19 36	-2	...	7.95	7.75	0.44	2.53	19.25	19.04	0.08	13.65	13.56	0.05	-20.58
A211615	...	...	11 48 30.7	20 25 29	-2	...	11.06	10.89	1.02	3.56	20.83	20.63	0.12	14.66	14.57	0.06	-19.57
U6851	N3937	127-088	11 50 07.9	20 54 38	-2	6618	11.71	11.55	0.17	3.77	18.68	18.47	0.03	10.95	10.85	0.07	-23.29
A211616	...	...	11 50 08.5	20 52 19	-2	...	5.85	5.60	0.48	1.83	19.06	18.83	0.11	14.48	14.38	0.05	-19.76
U6852	N3940	127-089	11 50 11.8	21 16 06	-5	6500	15.73	15.62	0.30	5.10	19.62	19.41	0.03	11.63	11.53	0.06	-22.62
A211617	...	...	11 50 14.8	20 56 53	-2	...	3.49	3.14	0.17	1.03	18.09	17.80	0.08	14.03	13.93	0.05	-20.21
A211618	...	...	11 50 27.4	20 57 51	-5	...	4.71	4.44	0.16	1.45	18.65	18.40	0.06	13.56	13.46	0.05	-20.68
A210878	...	127-092	11 50 36.3	20 56 13	-5	6918	9.24	9.05	0.24	2.96	19.74	19.52	0.04	12.93	12.83	0.06	-21.31
A211619	...	...	11 50 53.4	20 58 21	-2	...	24.21	24.15	1.69	7.89	21.33	21.13	0.08	14.62	14.52	0.06	-19.62
A211620	...	...	11 50 54.1	20 45 49	-2	...	10.84	10.67	0.76	3.48	21.12	20.91	0.09	14.77	14.67	0.06	-19.47
A221044	...	159-083	12 47 16.2	27 10 00	-5	7429	9.23	9.04	0.21	3.15	18.87	18.70	0.04	12.30	12.26	0.05	-22.02
A221740	...	...	12 47 25.3	27 10 35	-2	...	15.72	15.61	0.91	5.44	20.99	20.84	0.07	14.55	14.51	0.06	-19.77
U7986	N4715	159-085	12 47 31.6	28 05 40	1	6922	17.12	17.01	0.48	5.93	20.26	20.11	0.04	11.89	11.85	0.06	-22.43
A221739	...	...	12 47 47.0	27 02 52	-5	6980	8.08	7.88	0.38	2.74	19.77	19.61	0.07	13.39	13.35	0.06	-20.93
A221741	...	...	12 48 00.2	26 58 50	-2	5662	7.83	7.62	0.48	2.65	19.19	19.03	0.09	13.65	13.61	0.05	-20.67
A221057	...	159-089	12 48 26.6	28 06 33	-5	7468	7.59	7.38	0.20	2.57	19.08	18.91	0.04	12.72	12.68	0.05	-21.61
A221585	...	...	12 48 39.5	28 04 36	-2	7455	6.12	5.87	0.37	2.04	19.38	19.20	0.08	14.18	14.14	0.05	-20.15
A221742	...	...	12 50 14.9	26 38 12	-5	6295	8.23	8.07	0.38	2.81	19.87	19.71	0.07	13.44	13.40	0.06	-20.88
A221083	I831	159-100	12 50 17.7	26 44 29	-5	6366	13.82	13.72	0.31	4.78	19.60	19.45	0.04	12.32	12.28	0.06	-22.00
A221744	...	...	12 50 37.0	26 44 51	-5	...	4.28	4.04	0.21	1.41	19.16	18.97	0.07	14.44	14.40	0.06	-19.88
A221100	I832	159-105	12 51 30.0	26 43 18	-5	7005	7.95	7.78	0.18	2.71	19.00	18.84	0.04	12.59	12.55	0.05	-21.73
U8026	N4787	160-006	12 51 39.6	27 20 22	-5	7635	20.51	20.44	0.67	7.12	20.48	20.34	0.15	12.42	12.39	0.10	-21.89
U8028	N4789	160-008	12 51 53.1	27 20 25	-5	8372	16.43	16.34	0.22	5.69	19.05	18.91	0.15	11.14	11.11	0.10	-23.18
U8038	N4798	160-013	12 52 29.1	27 41 11	-2	7838	46.96	46.96	1.47	16.36	21.20	21.06	0.04	11.93	11.89	0.06	-22.39
U8049	N4807	160-017	12 53 03.8	27 47 28	-5	6941	7.05	6.80	0.18	2.37	18.34	18.17	0.04	12.29	12.25	0.05	-22.03
A221118	...	...	12 53 06.8	27 47 45	-2	7144	32.97	32.97	2.25	11.48	22.12	21.98	0.07	14.38	14.34	0.06	-19.94
A221121	I3900	160-019	12 53 15.6	27 31 37	-5	7171	5.36	5.08	0.23	1.77	17.89	17.71	0.07	12.63	12.60	0.05	-21.69
A221746	...	...	12 53 32.2	28 10 29	-5	6687	4.57	4.31	0.27	1.50	18.96	18.76	0.09	13.90	13.85	0.06	-20.43
A221123	...	160-020	12 53 40.5	27 56 50	-2	4802	4.34	4.08	0.41	1.42	18.70	18.51	0.13	14.38	14.35	0.06	-19.93

TABLE 3. (continued)

Name	NGC	CGCG	R.A. (1950)	Dec (1950)	T	V_{hel} (km/s)	r_e ($''$)	r_e^c ($''$)	ϵ_r ($''$)	R_e (kpc)	μ_e – (mag/arcsec ²) –	μ_e^c (13)	ϵ_μ (14)	m_I (mag)	m_I^c (mag)	ϵ_m (mag)	M_I (mag)
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)
A221124	...	...	12 53 40.8	27 55 06	-2	7419	47.86	47.86	3.57	16.67	22.18	22.03	0.07	14.63	14.59	0.06	-19.69
A221747	...	...	12 53 43.0	26 35 13	-2	...	43.86	43.86	6.14	15.28	22.63	22.49	0.13	15.20	15.17	0.05	-19.12
U8057	N4816	160-021	12 53 46.8	28 00 56	-2	6878	24.60	24.56	0.59	8.56	20.33	20.18	0.04	10.71	10.66	0.09	-23.62
A221748	...	...	12 53 49.3	27 46 35	-2	7451	11.48	11.36	0.60	3.96	21.00	20.85	0.07	15.14	15.10	0.07	-19.18
A221749	...	...	12 53 51.3	27 42 57	-5	...	6.13	5.92	0.34	2.06	19.68	19.51	0.08	13.90	13.86	0.05	-20.42
A221129	I834	160-022	12 53 52.7	26 37 45	-5	6292	10.39	10.23	0.32	3.56	19.19	19.04	0.05	12.39	12.36	0.06	-21.92
A221128	...	160-023	12 53 54.4	28 01 16	-5	7096	4.48	4.22	0.15	1.47	18.18	17.98	0.15	13.21	13.17	0.10	-21.12
A221750	...	...	12 54 01.1	27 59 49	-5	6836	2.15	1.67	0.13	0.58	17.62	17.24	0.09	14.39	14.34	0.05	-19.94
A221134	...	160-027	12 54 01.3	28 06 03	-5	6665	5.51	5.27	0.21	1.84	18.86	18.68	0.06	13.23	13.18	0.06	-21.10
U8060	N4819	160-025	12 54 02.8	27 15 30	1	6696	23.96	23.87	0.73	8.31	20.17	20.04	0.04	12.11	12.09	0.05	-22.20
A221132	...	D140	12 54 04.3	28 12 47	-5	6757	4.41	4.15	0.20	1.45	18.66	18.46	0.07	13.47	13.42	0.05	-20.86
A221133	N4821	160-024	12 54 05.1	27 14 00	-2	6974	11.87	11.69	0.36	4.07	19.44	19.30	0.04	13.00	12.98	0.05	-21.31
A221751	...	...	12 54 06.6	27 19 31	-5	6134	3.54	3.16	0.23	1.10	17.90	17.68	0.10	13.75	13.73	0.05	-20.56
A221135	N4824	...	12 54 08.3	27 48 40	-5	7105	4.77	4.54	0.20	1.58	18.76	18.57	0.06	13.48	13.44	0.05	-20.84
A221753	...	...	12 54 09.2	27 29 51	-2	7388	32.87	32.87	1.89	11.45	22.57	22.43	0.06	14.62	14.58	0.06	-19.70
U8065	N4827	160-028	12 54 18.0	27 26 55	-5	7650	12.61	12.45	0.26	4.34	19.18	19.04	0.04	11.81	11.78	0.05	-22.50
A221145	...	...	12 54 21.7	27 41 37	-2	7762	24.55	24.52	2.02	8.54	21.79	21.64	0.09	14.89	14.85	0.06	-19.43
A221147	...	160-031	12 54 24.3	27 21 48	-2	6849	82.29	82.29	4.94	28.66	22.64	22.51	0.06	13.69	13.67	0.06	-20.62
A221151	...	160-033	12 54 25.5	27 10 07	-2	6304	18.66	18.55	1.05	6.46	20.92	20.79	0.07	13.55	13.53	0.06	-20.75
A221154	...	D245	12 54 31.2	28 53 49	-2	...	21.67	21.58	2.32	7.52	20.87	20.72	0.12	13.69	13.64	0.06	-20.64
A221755	...	...	12 54 35.2	29 10 29	-5	...	3.37	3.08	0.21	1.07	18.97	18.74	0.09	14.59	14.55	0.05	-19.73
A221155	...	...	12 54 35.9	29 19 27	-5	7516	7.36	7.19	0.25	2.50	18.78	18.61	0.05	12.93	12.89	0.05	-21.40
A221752	...	...	12 54 36.4	27 38 31	-2	7012	18.89	18.81	0.97	6.55	21.21	21.07	0.06	14.35	14.31	0.05	-19.97
A221158	...	D34	12 54 38.4	27 47 52	-2	8328	62.93	62.93	5.41	21.92	23.20	23.05	0.08	15.31	15.26	0.05	-19.02
A221756	...	...	12 54 40.4	29 15 08	-2	...	4.04	3.79	0.27	1.32	18.82	18.62	0.10	14.48	14.44	0.05	-19.84
A221162	...	D16	12 54 44.6	27 40 39	-2	6215	16.51	16.43	0.84	5.72	21.20	21.06	0.06	13.84	13.80	0.06	-20.49
A221164	...	160-037	12 54 45.4	27 44 20	-5	7336	4.01	3.78	0.13	1.31	17.85	17.64	0.05	13.03	12.98	0.04	-21.31
U8069	...	160-038	12 54 46.5	29 19 19	-2	7460	20.94	20.88	1.19	7.27	19.89	19.74	0.07	12.31	12.27	0.05	-22.02
A221167	...	D94	12 54 52.7	28 04 57	-2	7084	30.91	30.91	2.19	10.77	21.85	21.69	0.08	14.73	14.68	0.06	-19.60
A221168	...	D112	12 54 56.3	28 09 10	-5	7428	4.55	4.29	0.26	1.49	19.24	19.03	0.08	14.25	14.20	0.06	-20.08
U8070	N4839	160-039	12 54 58.3	27 46 11	-2	7346	33.07	33.07	0.35	11.52	20.23	20.07	0.03	9.44	9.38	0.17	-24.90
A221172	...	160-040	12 55 00.6	27 48 57	-2	5517	4.90	4.71	0.18	1.64	18.43	18.24	0.06	13.76	13.71	0.05	-20.57
U8071	...	160-043	12 55 02.6	28 26 59	-2	7069	21.96	21.89	0.80	7.63	21.09	20.92	0.05	13.80	13.74	0.05	-20.54

TABLE 3. (continued)

Name	NGC	CGCG	R.A.	Dec	T	V_{hel}	r_e	r_e^c	ϵ_r	R_e	μ_e	μ_e^c	ϵ_μ	m_I	m_I^c	ϵ_m	M_I
(1)	(2)	(3)	(1950)	(1950)	(6)	(km/s)	($''$)	($''$)	($''$)	(kpc)	(mag/arcsec ²)	(mag/arcsec ²)	(14)	(mag)	(mag)	(mag)	(mag)
A221173	...	...	12 55 04.0	28 27 40	-2	...	70.10	70.10	2.55	24.42	22.02	21.85	0.04	13.15	13.09	0.07	-21.1
U8072	N4841A	160-044S	12 55 07.4	28 44 50	-5	6784	15.92	15.81	0.24	5.51	19.34	19.18	0.15	11.40	11.35	0.10	-22.1
A221175	N4840	160-042	12 55 07.5	27 52 55	-5	6059	6.81	6.66	0.15	2.32	18.43	18.25	0.04	12.43	12.38	0.05	-21.1
U8073	N4841B	160-044N	12 55 09.5	28 45 21	-5	6266	7.08	6.85	0.18	2.39	18.71	18.53	0.15	12.55	12.51	0.10	-21.1
A221178	N4842	160-046	12 55 10.2	27 45 25	-5	7297	5.06	4.87	0.09	1.70	18.10	17.91	0.03	12.75	12.69	0.05	-21.1
A221181	...	160-045	12 55 17.5	27 07 19	-2	6511	9.76	9.56	0.67	3.33	19.46	19.31	0.09	13.67	13.64	0.05	-20.1
A221757	...	...	12 55 22.3	29 25 13	-5	7165	4.76	4.53	0.24	1.58	19.03	18.84	0.08	13.83	13.79	0.05	-20.1
A221185	...	160-047	12 55 22.3	28 06 20	-2	6156	8.54	8.39	0.79	2.92	19.32	19.16	0.11	13.92	13.87	0.05	-20.1
A221186	...	160-048N	12 55 22.7	28 09 19	-5	5805	6.86	6.70	0.18	2.33	19.18	19.00	0.04	13.21	13.16	0.06	-21.1
A221188	...	160-049	12 55 22.9	28 27 12	-5	7350	3.01	2.65	0.10	0.92	17.61	17.34	0.05	13.61	13.55	0.05	-20.1
A221190	...	160-048S	12 55 25.5	28 09 05	-2	7046	6.57	6.40	0.36	2.23	19.70	19.53	0.07	13.85	13.80	0.06	-20.1
A221193	...	D238	12 55 28.7	28 46 21	-5	7338	3.02	2.70	0.14	0.94	18.07	17.83	0.07	14.06	14.01	0.05	-20.1
U8080	...	160-051	12 55 37.0	27 07 44	-2	7410	6.77	6.51	0.30	2.27	18.97	18.81	0.06	12.69	12.66	0.06	-21.1
A221200	...	160-052	12 55 37.7	27 11 08	-5	7976	7.09	6.84	0.33	2.38	19.40	19.23	0.07	13.39	13.36	0.06	-20.1
A221203	...	160-053	12 55 38.4	28 05 20	-2	7198	4.58	4.37	0.30	1.52	18.37	18.18	0.09	13.75	13.70	0.06	-20.1
A221758	...	...	12 55 46.6	28 12 33	-5	8509	2.97	2.69	0.20	0.94	19.24	19.00	0.10	15.06	15.01	0.06	-19.1
U8086	N4849	160-056S	12 55 47.5	26 40 01	-5	5885	13.17	13.04	0.22	4.54	19.12	18.97	0.03	11.75	11.72	0.06	-22.1
A221210	...	D209	12 55 48.3	28 27 20	-2	...	34.41	34.42	1.33	11.99	21.82	21.66	0.05	13.84	13.78	0.06	-20.1
A221208	...	160-059	12 55 48.7	29 13 04	-2	7560	67.34	67.34	3.86	23.46	21.60	21.45	0.06	13.23	13.18	0.05	-21.1
A221212	...	160-057	12 55 50.1	28 23 54	-2	7405	3.91	3.62	0.38	1.26	17.87	17.64	0.14	13.72	13.66	0.05	-20.1
A221216	...	160-062	12 55 53.6	29 23 53	-2	7837	26.53	26.52	1.34	9.24	21.23	21.08	0.06	13.36	13.31	0.06	-20.1
A221220	N4850	160-063	12 55 57.1	28 14 25	-5	5984	6.14	5.96	0.17	2.08	18.66	18.49	0.04	12.90	12.85	0.05	-21.1
A221222	...	...	12 56 03.5	28 47 59	-5	...	5.79	5.58	0.34	1.94	19.54	19.36	0.08	14.09	14.04	0.05	-20.1
A221223	...	D44	12 56 03.6	27 49 51	-2	7554	28.96	28.94	4.68	10.08	21.33	21.18	0.17	14.74	14.70	0.06	-19.1
A221226	...	160-065	12 56 05.4	28 17 13	-2	7246	7.39	7.23	0.19	2.52	19.01	18.83	0.04	12.77	12.72	0.05	-21.1
A221230	...	D182	12 56 07.2	28 19 19	-5	5897	2.10	1.62	0.11	0.56	17.70	17.31	0.08	14.52	14.47	0.05	-19.1
U8092	N4853	160-068	12 56 10.3	27 52 01	-2	7704	7.46	7.24	0.15	2.52	18.23	18.06	0.04	12.52	12.47	0.05	-21.1
A221234	I3943	160-069	12 56 11.4	28 23 08	-5	6818	14.54	14.43	0.58	5.03	19.82	19.65	0.05	12.46	12.41	0.07	-21.1
A221238	...	D28	12 56 12.7	27 48 55	-2	5962	5.12	4.86	0.31	1.69	19.12	18.93	0.09	14.47	14.43	0.05	-19.1
A221242	N4854	160-070	12 56 22.4	27 56 45	-2	8077	13.74	13.61	0.41	4.74	20.16	20.00	0.04	12.58	12.53	0.06	-21.1
A221244	I3946	160-210	12 56 23.9	28 04 53	-5	6101	7.08	6.89	0.24	2.40	18.59	18.42	0.05	12.78	12.73	0.05	-21.1
A221247	...	D181	12 56 26.3	28 21 21	-2	5932	4.57	4.31	0.42	1.50	18.84	18.64	0.13	14.03	13.98	0.05	-20.1
A221248	I3947	160-211	12 56 27.2	28 03 33	-2	5730	5.59	5.37	0.26	1.87	18.66	18.48	0.07	13.54	13.49	0.06	-20.1

TABLE 3. (continued)

Name	NGC	CGCG	R.A. (1950)	Dec (1950)	T	V_{hel} (km/s)	r_e ($''$)	r_e^c ($''$)	ϵ_r ($''$)	R_e (kpc)	μ_e – (mag/arcsec ²) –	μ_e^c (13)	ϵ_μ (14)	m_I (mag)	m_I^c (mag)	ϵ_m (mag)	M_I (mag)
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)
A221252	...	D136	12 56 30.6	28 14 12	-2	5652	1.76	1.33	0.10	0.46	16.89	16.47	0.09	14.34	14.29	0.05	-19.99
A221256	...	D135	12 56 35.7	28 14 23	-5	8312	4.94	4.72	0.27	1.64	19.62	19.43	0.08	14.42	14.36	0.05	-19.93
A221257	...	160-214	12 56 36.6	28 29 52	-2	8018	12.22	12.10	0.35	4.21	19.53	19.36	0.05	13.12	13.06	0.05	-21.23
U8097	N4859	160-071	12 56 37.0	27 05 06	-2	7055	12.59	12.43	0.37	4.33	19.49	19.35	0.04	12.25	12.22	0.06	-22.06
A221259	N4860	160-215	12 56 39.3	28 23 47	-5	7920	7.85	7.68	0.13	2.68	18.50	18.32	0.03	12.19	12.13	0.05	-22.15
A221265	...	D108	12 56 39.6	28 10 50	-2	6443	9.37	9.21	0.66	3.21	20.07	19.91	0.09	14.44	14.39	0.05	-19.89
A221261	...	D180	12 56 40.3	28 20 05	-5	6396	12.12	12.01	1.19	4.18	21.88	21.71	0.12	14.72	14.67	0.11	-19.61
A221263	I3955	160-216	12 56 41.1	28 16 06	-2	7895	11.32	11.20	0.43	3.90	19.94	19.77	0.05	12.83	12.77	0.06	-21.51
A221266	I3957	160-217	12 56 42.8	28 02 21	-5	6350	4.04	3.73	0.16	1.30	18.48	18.26	0.06	13.57	13.52	0.05	-20.76
A221267	I3960	160-219	12 56 43.1	28 07 35	-5	6709	5.07	4.85	0.20	1.69	18.65	18.47	0.06	13.28	13.24	0.05	-21.04
A221268	I3959	160-218	12 56 43.5	28 03 16	-5	7033	4.91	4.66	0.14	1.62	18.17	17.98	0.05	12.90	12.86	0.05	-21.43
A221274	N4864	160-221	12 56 48.4	28 14 54	-5	6789	9.12	8.96	0.16	3.12	18.99	18.82	0.03	12.37	12.32	0.05	-21.96
A221275	I3963	160-220	12 56 48.6	28 02 43	-5	6675	13.62	13.49	0.73	4.70	20.43	20.27	0.07	12.96	12.91	0.07	-21.37
A221276	...	160-223	12 56 49.3	28 20 53	-5	7799	16.49	16.40	0.58	5.71	21.13	20.97	0.05	13.05	13.00	0.06	-21.29
A221277	N4867	160-222	12 56 50.6	28 14 33	-5	4752	4.85	4.62	0.13	1.61	18.13	17.95	0.04	13.02	12.98	0.05	-21.31
U8100	N4865	160-224	12 56 54.9	28 21 14	-5	4643	9.08	8.92	0.17	3.11	18.39	18.23	0.15	12.11	12.07	0.10	-22.22
A221281	...	D107	12 56 55.6	28 09 25	-2	6439	12.74	12.63	0.42	4.40	20.51	20.35	0.05	14.33	14.28	0.06	-20.00
A221284	...	D106	12 56 58.1	28 10 05	-5	5092	3.76	3.50	0.18	1.22	18.67	18.46	0.15	14.07	14.03	0.10	-20.25
A221285	N4869	160-225	12 56 58.2	28 10 54	-5	6790	8.46	8.30	0.14	2.89	18.84	18.67	0.15	12.39	12.34	0.10	-21.94
A221289	...	D67	12 57 00.2	28 00 46	-5	6008	3.75	3.42	0.18	1.19	18.21	17.98	0.07	13.86	13.82	0.06	-20.46
A221290	...	D132	12 57 00.8	28 14 23	-5	7663	6.91	6.73	0.36	2.34	19.96	19.78	0.07	14.06	14.01	0.06	-20.27
A221291	...	D157	12 57 00.9	28 14 42	-5	6083	6.49	6.29	0.27	2.19	19.54	19.37	0.06	13.76	13.72	0.07	-20.57
A221293	...	D156	12 57 01.8	28 16 15	-2	6657	5.53	5.31	0.25	1.85	19.76	19.58	0.06	14.63	14.58	0.06	-19.70
A221295	...	D177	12 57 04.1	28 18 43	-2	6871	7.10	6.92	0.38	2.41	19.84	19.66	0.15	14.57	14.52	0.10	-19.76
A221296	I3967	160-226	12 57 04.7	28 07 15	-2	6950	6.44	6.24	0.33	2.17	18.64	18.46	0.07	13.48	13.44	0.06	-20.85
A221298	N4871	160-227	12 57 05.5	28 13 42	-2	7113	14.09	13.99	0.42	4.87	20.06	19.90	0.15	13.17	13.12	0.10	-21.16
A221299	...	D87	12 57 05.9	28 03 45	-5	7721	3.00	2.68	0.16	0.93	18.75	18.51	0.08	14.57	14.52	0.06	-19.76
A221301	I3973	160-228	12 57 06.3	28 09 20	-2	4745	4.05	3.80	0.15	1.32	17.85	17.65	0.06	12.95	12.91	0.05	-21.37
A221302	...	D176	12 57 06.9	28 19 05	-2	5579	7.80	7.63	0.44	2.66	19.25	19.08	0.15	13.82	13.77	0.10	-20.51
A221303	N4873	160-229	12 57 08.1	28 15 19	-2	5662	13.78	13.68	0.38	4.77	20.11	19.95	0.15	13.24	13.20	0.10	-21.08
A221304	N4872	160-230	12 57 09.6	28 13 09	-5	7177	3.55	3.27	0.12	1.14	17.76	17.54	0.15	13.30	13.25	0.10	-21.03
U8103	N4874	160-231	12 57 11.1	28 13 53	-5	7175	52.91	52.91	0.71	18.43	20.86	20.71	0.03	9.15	9.10	0.13	-25.18
A221313	N4875	160-232	12 57 13.2	28 10 43	-5	7897	4.48	4.25	0.15	1.48	18.37	18.17	0.15	13.40	13.35	0.10	-20.93

TABLE 3. (continued)

Name	NGC	CGCG	R.A. (1950)	Dec (1950)	T	V_{hel} (km/s)	r_e ($''$)	r_e^c ($''$)	ϵ_r ($''$)	R_e (kpc)	μ_e – (mag/arcsec ²) –	μ_e^c (13)	ϵ_μ (14)	m_I (mag)	m_I^c (mag)	ϵ_m (mag)	M_I (mag)
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)
A221314	...	D154	12 57 13.8	28 15 33	-2	6848	16.56	16.47	0.98	5.74	21.63	21.48	0.15	13.88	13.83	0.10	-20
A221317	...	D128	12 57 15.1	28 13 32	-2	7681	7.62	7.45	0.70	2.60	19.95	19.77	0.15	14.76	14.71	0.10	-19
A221320	...	D127	12 57 15.7	28 14 25	-2	7545	6.90	6.72	0.58	2.34	20.21	20.03	0.15	14.98	14.93	0.10	-19
A221323	...	160-233	12 57 17.9	28 11 47	-5	6899	1.35	0.91	0.08	0.32	16.51	15.93	0.15	14.35	14.30	0.10	-19
A221329	...	D153	12 57 19.2	28 15 57	-5	6652	2.98	2.66	0.15	0.93	18.18	17.94	0.15	14.03	13.98	0.10	-20
A221330	...	D126	12 57 19.6	28 13 49	-2	6932	12.77	12.66	0.67	4.41	20.66	20.50	0.15	14.66	14.61	0.10	-19
A221331	N4876	160-234	12 57 19.8	28 11 02	-2	6654	8.40	8.24	0.22	2.87	19.12	18.95	0.15	13.36	13.31	0.10	-20
A221332	...	160-235	12 57 21.5	28 07 42	-5	8028	7.79	7.62	0.37	2.65	19.54	19.36	0.07	13.47	13.42	0.06	-20
A221334	...	D152	12 57 22.5	28 14 46	-2	9371	11.42	11.30	0.40	3.94	20.10	19.93	0.15	13.56	13.50	0.10	-20
A221335	...	160-079	12 57 23.0	27 58 51	-2	8342	17.16	17.07	0.58	5.95	19.78	19.63	0.04	12.94	12.90	0.05	-21
A221343	...	D193	12 57 30.8	28 24 00	-2	7544	5.87	5.66	0.29	1.97	19.50	19.31	0.07	14.19	14.14	0.06	-20
A221345	N4883	160-237	12 57 31.5	28 18 24	-2	7961	8.99	8.83	0.25	3.08	19.34	19.17	0.04	12.94	12.89	0.06	-21
U8106	N4881	160-238	12 57 32.8	28 31 08	-5	6694	9.21	9.05	0.20	3.15	19.12	18.95	0.04	12.29	12.24	0.05	-22
A221354	N4886	160-239	12 57 40.0	28 15 32	-5	6345	9.37	9.21	0.28	3.21	19.59	19.43	0.15	12.78	12.73	0.10	-21
A221359	...	...	12 57 41.3	27 57 16	-5	7537	2.59	2.22	0.16	0.77	18.14	17.87	0.10	14.35	14.31	0.05	-19
A221357	...	D65	12 57 41.6	28 02 46	-5	6053	15.35	15.25	0.48	5.31	20.68	20.53	0.04	13.05	13.01	0.08	-21
A221362	I4011	160-242	12 57 42.0	28 16 33	-2	7216	5.85	5.64	0.34	1.96	19.36	19.18	0.15	14.03	13.98	0.10	-20
A221364	I4012	160-244	12 57 43.4	28 20 59	-2	7228	2.97	2.65	0.11	0.92	17.44	17.20	0.06	13.63	13.58	0.05	-20
U8110	N4889	160-241	12 57 43.7	28 14 54	-5	6497	35.74	35.74	0.33	12.45	19.61	19.46	0.03	9.45	9.40	0.09	-24
A221365	...	D207	12 57 43.8	28 26 31	-2	6730	7.46	7.29	0.24	2.54	19.40	19.22	0.05	13.66	13.61	0.06	-20
A221367	...	D40	12 57 45.8	27 51 53	-2	5589	12.87	12.76	0.67	4.44	21.34	21.18	0.06	14.97	14.93	0.06	-19
A221372	...	D173	12 57 48.4	28 20 48	-5	7384	4.51	4.28	0.23	1.49	18.71	18.52	0.08	13.84	13.79	0.05	-20
A221376	I4021	160-246	12 57 50.3	28 18 45	-5	5712	3.69	3.42	0.16	1.19	18.24	18.02	0.07	13.61	13.57	0.05	-20
A221377	N4894	160-247	12 57 52.1	28 14 20	-2	4587	14.25	14.15	0.85	4.93	20.50	20.36	0.15	14.01	13.97	0.10	-20
A221380	N4898A	160-248W	12 57 53.1	28 13 35	-2	6863	8.53	8.37	0.21	2.92	18.65	18.48	0.15	12.60	12.55	0.10	-21
U8113	N4895	160-249	12 57 53.5	28 28 15	-5	8406	24.41	24.38	0.60	8.49	19.77	19.61	0.04	11.34	11.28	0.07	-23
A221382	N4898B	160-248E	12 57 53.8	28 13 41	-2	6342	2.09	1.65	0.11	0.57	17.33	16.97	0.15	14.29	14.24	0.10	-20
A221385	...	D39	12 57 54.4	27 49 26	-2	5891	7.78	7.61	0.39	2.65	19.73	19.56	0.07	13.89	13.85	0.06	-20
A221386	I4026	160-250	12 57 57.7	28 19 04	-5	8189	8.24	8.08	0.28	2.81	19.61	19.43	0.05	13.24	13.19	0.06	-21
A221390	...	D27	12 58 01.6	27 47 11	-5	7460	6.15	5.94	0.27	2.07	19.56	19.38	0.06	13.81	13.77	0.06	-20
A221392	...	D119	12 58 03.5	28 13 38	-2	6927	5.65	5.43	0.28	1.96	19.43	19.26	0.15	14.21	13.88	0.10	-20
A221393	...	D147	12 58 03.8	28 14 36	-2	7744	22.77	22.73	1.04	7.92	21.40	21.24	0.15	13.76	13.71	0.10	-20
U8117	N4896	160-087	12 58 06.3	28 36 55	-2	5823	22.95	22.90	0.54	7.98	20.36	20.21	0.04	12.72	12.67	0.05	-21

TABLE 3. (continued)

Name	NGC	CGCG	R.A. (1950)	Dec (1950)	T	V_{hel} (km/s)	r_e ($''$)	r_e^c ($''$)	ϵ_r ($''$)	R_e (kpc)	μ_e – (mag/arcsec ²) –	μ_e^c (13)	ϵ_μ (14)	m_I (mag)	m_I^c (mag)	ϵ_m (mag)	M_I (mag)
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)
A221398	...	D63	12 58 07.7	28 02 10	-2	6707	14.56	14.46	1.07	5.04	21.14	20.99	0.09	14.66	14.61	0.06	-19.67
A221403	...	160-251	12 58 11.1	28 25 02	-2	5526	53.42	53.42	5.18	18.61	21.80	21.65	0.09	13.90	13.86	0.05	-20.42
A221404	...	D146	12 58 14.2	28 17 11	-2	7020	20.17	20.10	0.73	7.00	21.10	20.95	0.05	12.78	12.73	0.08	-21.55
A221408	N4906	160-253	12 58 15.2	28 11 42	-5	7477	7.36	7.19	0.19	2.50	19.04	18.86	0.04	12.87	12.82	0.05	-21.46
A221410	I4041	160-254	12 58 16.4	28 16 06	-2	7114	12.70	12.59	0.36	4.39	20.14	19.98	0.04	13.65	13.60	0.06	-20.68
A221412	I4042	160-255	12 58 18.3	28 14 32	-2	8046	6.29	6.09	0.24	2.12	18.67	18.48	0.06	12.84	12.79	0.05	-21.49
A221413	...	D116	12 58 18.5	28 14 04	-5	6255	7.08	6.90	0.29	2.40	19.54	19.37	0.06	13.50	13.46	0.06	-20.83
A221415	...	160-089	12 58 19.3	28 36 31	-2	7997	5.64	5.41	0.25	1.88	18.47	18.28	0.06	13.25	13.19	0.06	-21.09
A221416	...	D191	12 58 20.4	28 22 08	-2	6611	15.67	15.57	1.81	5.42	20.70	20.55	0.13	14.52	14.47	0.05	-19.81
A221419	I4045	160-256	12 58 24.5	28 21 40	-2	6926	7.52	7.35	0.18	2.56	18.52	18.35	0.04	12.80	12.75	0.06	-21.53
A221420	N4908	160-258	12 58 27.2	28 18 48	-5	8838	8.48	8.32	0.20	2.90	18.75	18.57	0.04	12.39	12.34	0.06	-21.95
A221422	...	160-091	12 58 28.2	28 38 00	-2	7684	10.93	10.78	0.27	3.76	19.32	19.15	0.04	12.35	12.30	0.06	-21.99
U8129	I4051	160-259	12 58 30.1	28 16 42	-5	4964	21.01	20.95	0.40	7.30	20.26	20.11	0.03	11.84	11.80	0.06	-22.48
A221429	...	160-261	12 58 35.0	28 10 13	-2	6907	21.38	21.32	1.05	7.43	20.61	20.46	0.06	13.62	13.57	0.06	-20.71
A221433	...	D229	12 58 44.2	28 37 53	-5	...	6.91	6.70	0.41	2.33	19.99	19.81	0.08	13.94	13.89	0.06	-20.39
A221434	...	160-092	12 58 44.7	28 05 16	-2	5977	6.56	6.36	0.32	2.21	19.65	19.47	0.07	13.76	13.72	0.05	-20.57
A221435	...	160-093	12 58 46.4	28 04 20	-5	6513	8.37	8.19	0.52	2.85	19.90	19.73	0.08	13.50	13.45	0.08	-20.83
U8133	N4919	160-094	12 58 53.2	28 04 44	-2	7110	9.51	9.34	0.23	3.25	18.97	18.79	0.04	12.48	12.43	0.06	-21.85
A221438	...	D204	12 58 57.8	28 27 59	-2	7667	5.28	5.05	0.21	1.76	19.08	18.89	0.06	13.74	13.69	0.06	-20.60
A221442	N4923	160-097	12 59 07.4	28 07 02	-2	5458	6.43	6.21	0.12	2.16	18.38	18.20	0.04	12.16	12.11	0.06	-22.17
U8137	I843	160-099	12 59 09.5	29 23 58	-5	7503	15.88	15.77	0.50	5.49	19.34	19.19	0.05	11.88	11.85	0.06	-22.43
A221599	...	...	12 59 19.9	29 16 05	-2	6386	50.89	50.89	2.18	17.73	22.06	21.92	0.05	14.10	14.07	0.06	-20.21
A221449	...	160-101	12 59 23.2	28 21 57	-5	5917	11.34	11.22	0.73	3.91	19.30	19.13	0.08	12.50	12.45	0.07	-21.83
A221450	...	D36	12 59 23.5	27 52 26	-2	8223	27.07	27.05	1.85	9.42	21.58	21.43	0.07	14.44	14.40	0.05	-19.88
A221453	...	160-100	12 59 26.2	28 09 47	-2	7616	3.56	3.27	0.15	1.14	18.20	17.96	0.07	13.25	13.20	0.06	-21.10
U8142	N4926	160-103	12 59 29.7	27 53 35	-2	7888	9.15	8.98	0.15	3.13	18.47	18.30	0.03	11.74	11.69	0.05	-22.59
A221455	N4927	160-105	12 59 33.6	28 16 35	-5	7583	11.58	11.46	0.28	3.99	19.17	19.00	0.04	12.12	12.06	0.06	-22.22
A221460	...	160-107	12 59 40.1	29 31 44	-2	7367	28.17	28.15	2.37	9.81	19.93	19.79	0.09	12.72	12.69	0.05	-21.59
A221463	...	160-109	12 59 44.1	28 39 27	-5	7613	10.58	10.40	0.38	3.62	19.57	19.40	0.05	12.72	12.67	0.06	-21.61
A221470	...	160-110	12 59 56.9	28 30 08	-2	5721	17.13	17.03	0.94	5.93	20.06	19.92	0.06	13.49	13.45	0.05	-20.84
A221471	...	160-111	12 59 57.4	28 31 28	-2	7137	19.79	19.69	0.74	6.86	20.78	20.63	0.05	13.77	13.73	0.06	-20.56
A230007	...	D113	13 00 10.5	28 13 09	-2	5779	47.43	47.43	2.09	16.52	22.51	22.35	0.05	14.06	14.00	0.06	-20.28
A230012	...	160-114	13 00 17.2	28 38 41	-2	7441	39.81	39.81	3.53	13.87	21.22	21.07	0.09	13.41	13.37	0.05	-20.92

TABLE 3. (continued)

Name	NGC	CGCG	R.A. (1950)	Dec (1950)	T	V_{hel} (km/s)	r_e ($''$)	r_e^c ($''$)	ϵ_r ($''$)	R_e (kpc)	μ_e – (mag/arcsec ²) –	μ_e^c (13)	ϵ_μ (14)	m_I (mag)	m_I^c (mag)	ϵ_m (mag)	M_I (mag)
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)
A230015	N4929	160-113	13 00 20.5	28 18 54	-2	6342	9.23	9.09	0.21	3.17	19.45	19.28	0.04	12.69	12.64	0.06	-21.65
A230022	I4111	160-116	13 00 32.7	28 20 24	-2	7893	26.47	26.47	0.99	9.22	21.39	21.22	0.05	13.90	13.84	0.06	-20.44
U8154	N4931	160-118	13 00 37.0	28 18 07	-5	5849	11.42	11.32	0.45	3.94	18.46	18.29	0.06	11.86	11.81	0.05	-22.47
A230030	...	160-119	13 00 45.7	28 51 06	-2	6917	11.94	11.79	0.97	4.11	19.36	19.20	0.10	13.12	13.08	0.06	-21.20
U8160	N4934	160-120	13 00 52.1	28 17 54	0	6041	248.2	248.28	25.87	86.49	22.53	22.38	0.07	13.18	13.13	0.05	-21.15
A230038	N4943	160-122	13 01 20.8	28 21 08	-2	5739	15.71	15.61	0.68	5.44	20.31	20.16	0.06	13.24	13.20	0.06	-21.09
U8167	N4944	160-124	13 01 25.8	28 27 24	-2	7111	33.48	33.49	0.59	11.67	19.92	19.77	0.03	11.86	11.82	0.05	-22.47
A230042	I4133	160-123	13 01 26.6	28 15 23	-2	6330	3.93	3.65	0.16	1.27	18.14	17.91	0.06	13.22	13.17	0.05	-21.11
A231125	...	...	13 01 43.7	28 14 00	-2	6123	22.02	21.95	1.35	7.65	21.72	21.56	0.07	14.17	14.12	0.06	-20.16
A230046	...	160-125	13 01 46.2	28 30 57	-5	5971	7.10	6.90	0.31	2.40	19.27	19.10	0.06	13.15	13.11	0.06	-21.17
A231126	...	...	13 01 48.8	28 27 44	-5	...	2.96	2.59	0.26	0.90	17.04	16.79	0.13	13.34	13.30	0.05	-20.98
U8175	N4952	160-129	13 02 35.4	29 23 28	-5	5919	12.90	12.78	0.20	4.45	18.81	18.65	0.03	11.53	11.49	0.05	-22.79
A231127	...	...	13 02 44.5	27 46 48	-2	...	49.37	49.37	2.73	17.20	22.66	22.51	0.06	14.85	14.81	0.06	-19.47
U8178	N4957	160-130	13 02 48.4	27 50 15	-5	6876	17.66	17.56	0.29	6.12	19.70	19.55	0.03	11.64	11.60	0.06	-22.68
A230071	...	160-135	13 03 41.9	28 51 36	-5	7994	5.01	4.75	0.20	1.65	18.44	18.25	0.06	13.19	13.14	0.05	-21.14
A231130	...	...	13 04 13.6	28 36 18	-2	...	17.75	17.64	0.90	6.14	21.67	21.52	0.06	14.48	14.44	0.06	-19.84
A230086	N4971	160-140	13 04 30.9	28 48 46	-5	6399	7.81	7.60	0.27	2.65	18.92	18.75	0.05	12.60	12.56	0.06	-21.72
A231131	...	...	13 04 32.9	28 46 17	-2	...	7.32	7.10	0.84	2.47	20.68	20.51	0.15	15.39	15.35	0.06	-18.93
U8260	N5004	160-157	13 08 39.3	29 54 08	-5	7141	11.76	11.60	0.19	4.04	18.81	18.66	0.03	11.65	11.61	0.05	-22.67
A330131	...	...	23 11 47.5	9 39 54	0	3894	5.95	5.74	0.34	0.98	19.54	19.40	0.08	13.93	13.87	0.05	-18.86
A330144	...	406-031	23 12 16.3	7 26 44	1	4714	300.2	300.23	20.96	51.22	24.51	24.38	0.05	14.46	14.38	0.05	-18.35
U12454	...	406-033	23 12 37.7	9 24 27	-5	4793	34.61	34.61	0.72	5.90	20.62	20.50	0.03	11.49	11.43	0.07	-21.30
A331631	...	...	23 12 39.5	9 23 36	-2	...	3.50	3.22	0.22	0.55	18.42	18.24	0.15	13.98	13.92	0.10	-18.81
A331632	...	...	23 12 44.1	9 21 29	-5	...	3.37	3.08	0.29	0.53	19.34	19.15	0.13	15.10	15.04	0.05	-17.69
A331633	...	...	23 12 53.1	7 33 31	-2	...	24.69	24.67	2.39	4.21	22.15	22.02	0.10	15.48	15.40	0.05	-17.33
A331634	...	...	23 13 05.5	7 35 43	-5	...	3.14	2.83	0.38	0.48	18.93	18.71	0.18	14.76	14.68	0.05	-18.05
A330161	N7557	406-035	23 13 07.8	6 26 00	-2	3754	20.59	20.51	0.61	3.50	20.61	20.44	0.04	12.84	12.72	0.05	-20.01
A331635	...	...	23 13 08.3	7 37 34	0	...	16.33	16.23	0.87	2.77	21.17	21.04	0.07	13.22	13.14	0.06	-19.59
U12464	N7562	406-039	23 13 25.1	6 24 53	-5	3514	27.36	27.36	0.27	4.67	18.98	18.81	0.03	10.05	9.93	0.05	-22.80
A331636	...	...	23 13 27.1	6 21 22	-2	...	41.58	41.58	2.28	7.09	22.40	22.23	0.06	14.38	14.26	0.05	-18.47
U12472	...	406-041	23 14 13.5	8 37 58	9	6358	10.18	10.08	0.36	1.72	18.78	18.62	0.05	12.04	11.95	0.05	-20.79
A331638	...	...	23 16 47.1	7 49 39	-5	...	6.69	6.48	0.36	1.11	19.94	19.76	0.08	13.90	13.81	0.05	-18.93

TABLE 3. (continued)

Name	NGC	CGCG	R.A. (1950)	Dec (1950)	T	V_{hel} (km/s)	r_e ($''$)	r_e^c ($''$)	ϵ_r ($''$)	R_e (kpc)	μ_e – (mag/arcsec ²) –	μ_e^c (13)	ϵ_μ (14)	m_I (mag)	m_I^c (mag)	ϵ_m (mag)	M_I (mag)
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)
A331639	...	...	23 16 47.9	8 11 37	-2	...	9.64	9.49	1.47	1.62	21.39	21.24	0.19	15.91	15.83	0.05	-16.90
A331640	...	...	23 17 03.3	8 20 14	-5	...	3.23	2.93	0.19	0.50	18.33	18.12	0.09	14.00	13.91	0.05	-18.82
U12509	N7611	406-066	23 17 04.9	7 47 24	-2	3252	5.72	5.51	0.14	0.94	17.08	16.91	0.04	11.35	11.26	0.05	-21.47
U12510	...	...	23 17 07.6	7 59 23	-2	3542	40.11	40.11	1.84	6.84	22.32	22.18	0.05	12.85	12.76	0.07	-19.97
U12512	N7612	406-068	23 17 12.2	8 18 09	-2	3228	13.10	13.00	0.27	2.22	18.55	18.40	0.04	11.21	11.13	0.05	-21.61
A331641	...	...	23 17 15.2	7 58 60	-5	...	3.78	3.49	0.35	0.60	19.90	19.69	0.13	15.20	15.11	0.05	-17.62
A331642	...	...	23 17 15.4	8 19 44	-2	...	4.45	4.21	0.27	0.72	19.66	19.48	0.08	15.19	15.10	0.05	-17.63
A331643	...	...	23 17 17.8	7 55 02	-2	...	10.69	10.53	0.67	1.80	20.95	20.80	0.08	14.73	14.64	0.05	-18.09
A330238	...	406-069	23 17 25.0	8 17 12	1	3429	122.8	122.84	7.44	20.96	23.13	22.99	0.05	13.28	13.20	0.06	-19.54
A330246	N7617	406-072	23 17 37.0	7 53 31	-2	4072	7.51	7.32	0.16	1.25	18.36	18.20	0.15	12.36	12.27	0.10	-20.46
A331644	...	...	23 17 37.1	7 26 55	-2	...	154.4	154.49	10.44	26.36	23.43	23.28	0.05	14.21	14.12	0.06	-18.62
U12523	N7619	406-073	23 17 42.6	7 55 57	-2	3758	20.57	20.50	0.21	3.50	18.40	18.26	0.03	9.59	9.50	0.06	-23.23
A330250	N7621	406-074	23 17 48.0	8 04 00	-2	...	31.49	31.49	1.08	5.37	21.35	21.21	0.03	13.82	13.73	0.05	-19.00
A331645	...	...	23 17 50.5	7 15 49	-5	...	6.11	5.90	0.20	1.01	19.04	18.84	0.05	13.29	13.18	0.06	-19.56
U12526	N7623	406-075	23 17 58.0	8 07 20	-2	3739	12.84	12.73	0.23	2.17	18.50	18.35	0.03	11.24	11.15	0.05	-21.58
U12531	N7626	406-076	23 18 10.3	7 56 35	-5	3416	27.78	27.78	0.32	4.74	19.15	19.01	0.03	9.88	9.79	0.05	-22.94
A331646	...	...	23 18 15.5	7 58 24	-2	...	2.12	1.64	0.41	0.28	18.41	18.03	0.15	15.74	15.65	0.10	-17.08
A331647	...	...	23 18 19.3	7 36 41	-5	...	5.87	5.66	0.73	0.97	20.41	20.24	0.16	15.13	15.03	0.06	-17.70
A330257	...	...	23 18 24.0	7 54 45	-2	4321	4.00	3.72	0.32	0.63	19.79	19.59	0.11	14.90	14.81	0.05	-17.92
A330258	...	...	23 18 28.5	7 38 37	-2	3797	95.75	95.75	4.93	16.34	22.68	22.53	0.05	14.07	13.98	0.06	-18.76
A331566	...	...	23 18 34.2	7 54 00	-5	...	11.26	11.13	0.27	1.90	19.88	19.73	0.04	13.04	12.95	0.05	-19.78
A330266	...	406-078	23 18 41.0	7 05 30	0	3076	10.85	10.72	0.19	1.83	18.76	18.59	0.03	11.80	11.69	0.06	-21.04
A331648	...	...	23 18 44.0	7 34 50	-5	...	20.13	20.06	1.44	3.42	21.21	21.06	0.08	13.03	12.94	0.14	-19.80
A331649	...	...	23 18 59.1	7 09 06	-5	...	3.52	3.24	0.33	0.55	19.45	19.21	0.13	14.98	14.87	0.06	-17.87
A331650	...	...	23 19 06.3	8 39 37	-5	...	1.73	1.26	0.11	0.21	17.39	16.95	0.10	14.60	14.53	0.06	-18.20
U12542	N7634	406-085	23 19 10.0	8 36 47	-5	3214	14.64	14.54	0.29	2.48	18.95	18.83	0.03	11.46	11.39	0.06	-21.34
A331651	...	...	23 19 43.9	8 11 12	-5	...	5.44	5.22	0.35	0.89	19.97	19.82	0.09	14.57	14.50	0.05	-18.23
A331652	...	...	23 20 56.5	9 28 09	-5	...	15.29	15.19	0.99	2.59	21.61	21.50	0.07	14.90	13.79	0.08	-19.00
A331653	...	...	23 21 17.1	9 28 11	-5	...	6.70	6.51	0.34	1.11	19.51	19.38	0.07	13.68	13.62	0.05	-19.11
U12575	N7648	406-096	23 21 22.2	9 23 37	1	3559	40.69	40.69	1.47	6.94	20.75	20.64	0.05	11.76	11.70	0.05	-21.03
A330499	...	476-052	23 27 34.8	25 44 14	-5	9454	5.89	5.70	0.18	2.46	18.93	18.67	0.05	13.36	13.26	0.04	-21.49
A331654	...	...	23 28 33.2	25 48 35	-5	...	5.81	5.62	0.34	2.42	20.07	19.81	0.08	14.39	14.29	0.05	-20.46

TABLE 3. (continued)

Name	NGC	CGCG	R.A. (1950)	Dec (1950)	T	V_{hel} (km/s)	r_e ($''$)	r_e^c ($''$)	ϵ_r ($''$)	R_e (kpc)	μ_e – (mag/arcsec ²)	μ_e^c – (mag/arcsec ²)	ϵ_μ – (mag/arcsec ²)	m_I (mag)	m_I^c (mag)	ϵ_m (mag)	M_I (mag)
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)
A331655	...	...	23 28 59.1	25 51 41	-2	...	1.70	1.27	0.12	0.55	16.45	15.93	0.11	14.24	14.14	0.04	-20.61
A331656	...	...	23 29 51.7	25 37 39	-2	...	2.63	2.31	0.17	1.00	18.42	18.10	0.10	14.90	14.81	0.05	-19.93
A331657	...	...	23 30 28.4	25 41 05	-2	...	9.24	9.10	0.32	3.92	19.94	19.71	0.05	13.21	13.12	0.05	-21.62
A331672	...	...	23 30 27.8	25 37 47	-5	...	2.89	2.60	0.23	1.12	18.81	18.50	0.12	15.01	14.92	0.04	-19.83
A331658	...	...	23 31 35.5	26 38 02	-5	...	2.56	2.23	0.18	0.96	18.93	18.59	0.11	15.18	15.08	0.06	-19.67
A331209	...	...	23 32 01.7	26 50 53	-2	9585	37.55	37.55	2.04	16.19	22.41	22.18	0.06	14.22	14.12	0.05	-20.63
A330553	...	476-071	23 32 08.9	26 26 49	-2	9700	10.20	10.08	0.35	4.35	19.52	19.27	0.05	13.10	12.99	0.06	-21.75
A331210	...	...	23 32 10.3	27 03 15	-5	8893	4.89	4.66	0.31	2.01	19.36	19.10	0.09	14.46	14.36	0.05	-20.38
A331222	...	...	23 33 12.5	26 16 58	-2	8639	16.37	16.27	0.79	7.02	21.81	21.59	0.06	14.92	14.83	0.07	-19.92
A331223	...	...	23 33 15.8	26 19 44	-2	9511	8.18	8.01	0.44	3.45	19.99	19.74	0.07	14.29	14.19	0.06	-20.55
A331659	...	...	23 33 32.9	26 10 45	-5	...	3.75	3.49	0.23	1.51	19.49	19.20	0.09	14.82	14.72	0.06	-20.02
A331231	...	...	23 33 56.0	27 07 17	-2	9247	13.60	13.50	0.53	5.82	20.12	19.88	0.05	13.67	13.57	0.06	-21.18
A331233	...	...	23 33 59.9	26 07 05	-5	9064	6.47	6.27	0.40	2.70	19.84	19.59	0.08	13.99	13.89	0.07	-20.85
A330574	...	...	23 34 08.8	26 37 58	-5	8934	3.18	2.88	0.23	1.24	19.07	18.75	0.11	14.93	14.82	0.06	-19.92
A330575	...	...	23 34 08.8	27 01 46	-2	8415	6.13	5.92	0.43	2.55	20.57	20.31	0.09	15.73	15.63	0.06	-19.12
A330590	...	...	23 34 39.3	26 41 25	-2	9477	30.51	30.51	4.06	13.16	21.53	21.29	0.13	14.68	14.57	0.06	-20.18
A330594	...	...	23 34 49.8	26 52 13	-2	9217	5.30	5.08	0.36	2.19	20.15	19.88	0.09	15.32	15.21	0.06	-19.54
A330595	...	476-079	23 34 49.9	26 36 04	-2	8335	10.47	10.33	0.44	4.45	19.56	19.31	0.06	13.34	13.23	0.07	-21.51
A330600	...	476-080	23 34 55.8	26 47 40	-2	10072	7.49	7.32	0.32	3.16	19.74	19.48	0.06	13.98	14.87	0.06	-19.88
A330602	...	...	23 34 57.5	26 21 21	-2	9650	8.11	7.94	0.45	3.42	19.65	19.40	0.07	14.19	14.09	0.06	-20.66
A330604	...	...	23 35 01.3	26 33 37	-5	8253	2.66	2.30	0.16	0.99	18.60	18.25	0.09	14.80	14.69	0.06	-20.05
A330614	...	476-081	23 35 14.7	26 13 41	-2	8682	30.79	30.80	1.72	13.28	21.05	20.82	0.06	13.52	13.42	0.06	-21.32
A330615	...	...	23 35 16.4	26 42 14	-5	11085	6.15	5.94	0.30	2.56	19.35	19.07	0.07	13.83	13.71	0.07	-21.03
A330621	...	...	23 35 25.1	26 49 24	-5	10187	6.44	6.28	0.34	2.71	20.13	19.87	0.07	14.29	14.18	0.05	-20.58
A330624	...	...	23 35 27.6	26 59 14	-5	10204	7.13	6.93	0.31	2.99	19.35	19.09	0.06	13.27	13.16	0.06	-21.59
A330631	...	...	23 35 31.9	26 55 31	-5	7956	3.02	2.76	0.26	1.19	19.41	19.11	0.12	15.28	15.18	0.05	-19.57
A330637	...	...	23 35 39.9	26 45 12	-2	8378	13.24	13.16	0.88	5.67	21.42	21.18	0.08	15.37	15.26	0.05	-19.49
A331239	...	...	23 35 42.1	26 13 02	-5	9319	6.54	6.35	0.32	2.74	19.18	18.93	0.07	13.58	13.48	0.07	-21.27
A330643	...	...	23 35 43.4	27 08 13	-2	9285	5.80	5.57	0.24	2.40	18.49	18.22	0.06	13.46	13.35	0.06	-21.40
A330642	...	...	23 35 43.9	26 41 25	-5	9310	3.50	3.22	0.27	1.39	19.58	19.27	0.11	15.47	15.36	0.07	-19.39
A330647	...	476-084	23 35 45.7	26 32 25	-2	8656	8.36	8.22	0.84	3.54	19.48	19.24	0.13	14.10	14.00	0.06	-20.75
A330648	...	...	23 35 47.5	26 45 33	-5	9552	2.28	2.04	0.12	0.88	18.02	17.69	0.15	14.43	14.32	0.10	-20.43
A330649	...	476-085	23 35 48.1	26 36 35	-5	9244	4.92	4.75	0.15	2.05	18.66	18.40	0.05	13.30	13.20	0.05	-21.56

TABLE 3. (continued)

Name	NGC	CGCG	R.A.	Dec	T	V_{hel}	r_e	r_e^c	ϵ_r	R_e	μ_e	μ_e^c	ϵ_μ	m_I	m_I^c	ϵ_m	M_I
(1)	(2)	(3)	(1950)	(1950)	(6)	(km/s)	($''$)	($''$)	($''$)	(kpc)	(mag/arcsec ²)	(mag/arcsec ²)	(14)	(mag)	(mag)	(mag)	(mag)
A330651	...	...	23 35 50.3	26 46 26	-5	10111	5.32	5.19	0.23	2.24	19.36	19.10	0.15	14.05	13.94	0.10	-20.
A330652	...	...	23 35 50.6	27 01 37	-5	9871	4.46	4.20	0.31	1.81	19.40	19.11	0.10	14.42	14.31	0.06	-20.
A330653	...	476-086	23 35 52.4	26 52 52	-5	9280	5.69	5.51	0.14	2.38	18.76	18.50	0.04	13.23	13.12	0.06	-21.
A330658	I5341	476-087	23 35 56.5	26 42 29	-5	10820	7.31	7.20	0.19	3.10	19.39	19.14	0.15	13.01	12.90	0.10	-21.
A330659	...	...	23 35 56.8	26 48 48	-2	7532	33.11	33.11	3.11	14.28	21.70	21.47	0.09	14.83	14.73	0.04	-20.
A330662	...	476-089	23 35 58.2	26 32 29	-5	9542	5.80	5.63	0.17	2.43	18.47	18.21	0.05	13.06	12.95	0.06	-21.
A330660	...	476-090	23 35 58.8	26 42 06	-5	9600	8.24	8.14	0.22	3.51	19.38	19.13	0.15	12.87	12.76	0.10	-21.
U12716	N7720A	476-091S	23 35 59.1	26 45 14	-5	9141	18.43	18.41	0.18	7.94	19.58	19.34	0.03	10.14	10.03	0.09	-24.
A331456	N7720B	476-091N	23 35 59.2	26 45 26	-5	8049	3.36	3.18	0.11	1.37	17.86	17.58	0.15	13.81	13.70	0.10	-21.
A330665	...	...	23 36 02.9	26 45 28	-2	9828	3.71	3.54	0.16	1.53	18.68	18.41	0.15	14.17	14.06	0.10	-20.
A330667	...	476-092	23 36 04.1	26 42 09	-5	8515	1.89	1.60	0.08	0.69	16.80	16.43	0.07	13.93	13.82	0.04	-20.
A330668	...	476-090B	23 36 05.9	26 45 09	-2	8378	13.74	13.67	0.59	5.90	20.16	19.92	0.15	13.58	13.47	0.10	-21.
A330678	...	...	23 36 07.9	26 44 00	-5	9356	5.82	5.69	0.13	2.45	18.79	18.53	0.15	13.06	12.95	0.10	-21.
A330679	...	476-095B	23 36 11.3	26 56 17	-2	9415	9.23	9.06	0.57	3.91	20.59	20.33	0.08	15.12	15.01	0.06	-19.
A330680	...	...	23 36 11.7	26 50 29	-5	8697	2.75	2.55	0.12	1.10	18.08	17.77	0.07	14.38	14.27	0.04	-20.
A330683	...	...	23 36 12.7	26 45 53	-5	9195	3.73	3.56	0.27	1.54	19.49	19.22	0.10	14.92	14.81	0.05	-19.
A330686	...	476-095A	23 36 13.3	26 56 17	-5	9201	14.78	14.67	0.55	6.33	19.87	19.62	0.05	12.41	12.30	0.07	-22.
A330687	...	...	23 36 13.8	26 34 25	-2	9112	7.80	7.68	0.37	3.31	19.22	18.98	0.07	13.32	13.22	0.05	-21.
A330688	...	...	23 36 15.9	26 53 43	-2	9934	10.78	10.66	0.83	4.60	20.27	20.03	0.09	14.42	14.31	0.05	-20.
A330691	...	...	23 36 18.9	26 50 45	-2	9066	5.22	5.08	0.64	2.19	19.04	18.78	0.16	14.54	14.43	0.04	-20.
A330695	...	...	23 36 20.2	26 44 19	-2	10023	7.95	7.85	0.48	3.39	20.35	20.10	0.08	14.66	17.88	0.05	-16.
A330696	...	476-096	23 36 20.3	26 59 26	-5	9297	8.89	8.69	0.16	3.75	18.86	18.60	0.03	12.38	12.27	0.06	-22.
A330698	...	...	23 36 22.7	26 47 43	-2	...	6.15	6.03	0.51	2.60	20.15	19.89	0.11	15.38	15.27	0.05	-19.
A330699	...	...	23 36 23.5	26 46 08	-5	9643	1.10	0.76	0.08	0.33	16.39	15.75	0.11	14.87	14.76	0.04	-19.
A330700	...	...	23 36 25.8	26 53 04	-5	9276	3.76	3.53	0.16	1.52	18.42	18.13	0.07	14.06	13.95	0.06	-20.
A330702	...	...	23 36 28.4	26 58 41	-2	8666	6.48	6.26	0.54	2.70	20.29	20.02	0.11	15.59	15.48	0.06	-19.
A330703	...	...	23 36 31.0	26 40 54	-5	9056	2.37	2.14	0.17	0.92	18.51	18.19	0.11	15.14	15.03	0.05	-19.
A330704	...	...	23 36 31.1	27 01 22	0	9752	21.63	21.54	1.54	9.29	21.80	21.56	0.08	14.37	14.26	0.06	-20.
A331246	...	...	23 36 31.2	26 17 19	-5	10542	3.62	3.40	0.16	1.47	18.68	18.39	0.07	14.11	14.00	0.06	-20.
A330706	...	...	23 36 32.2	26 49 31	-5	9890	4.48	4.32	0.16	1.86	18.75	18.48	0.15	13.77	13.66	0.10	-21.
A330711	...	...	23 36 36.6	26 55 43	-5	9702	3.60	3.43	0.21	1.48	19.08	18.81	0.08	14.63	14.52	0.05	-20.
A331660	...	...	23 36 36.6	27 16 07	-2	...	11.22	11.02	1.15	4.75	21.32	21.07	0.13	15.43	15.33	0.07	-19.
A331466	...	...	23 36 38.2	27 07 45	-2	8696	14.03	13.87	1.26	5.98	21.33	21.09	0.11	15.00	14.90	0.06	-19.

TABLE 3. (continued)

Name	NGC	CGCG	R.A. (1950)	Dec (1950)	T	V_{hel} (km/s)	r_e ($''$)	r_e^c ($''$)	ϵ_r ($''$)	R_e (kpc)	μ_e – (mag/arcsec ²) –	μ_e^c (13)	ϵ_μ (14)	m_I (mag)	m_I^c (mag)	ϵ_m (mag)	M_I (mag)
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)
A331248	...	...	23 36 41.9	27 16 25	-2	9314	8.88	8.66	0.39	3.73	19.66	19.40	0.06	13.37	13.26	0.06	-21.48
A330730	...	...	23 36 50.1	26 28 32	-5	9424	5.75	5.58	0.25	2.41	19.44	19.18	0.06	13.96	13.85	0.06	-20.89
A330731	...	...	23 36 58.0	27 01 11	-2	9310	8.06	7.81	0.71	3.37	20.83	20.57	0.12	15.34	15.23	0.06	-19.51
A330735	...	...	23 37 03.5	27 03 14	-2	10138	4.25	3.89	0.59	1.68	20.22	19.90	0.20	15.70	15.59	0.06	-19.15
A330741	...	476-101	23 37 18.9	27 05 56	1	10023	6.20	5.92	0.20	2.55	18.86	18.59	0.05	12.92	12.81	0.07	-21.93
A331481	...	...	23 37 24.5	26 23 50	-5	8686	8.31	7.97	0.78	3.44	20.71	20.45	0.13	14.20	14.10	0.07	-20.65
A330744	...	476-102	23 37 28.0	26 33 25	-5	8659	11.24	10.96	0.52	4.73	19.78	19.53	0.15	12.81	12.71	0.10	-22.04
U12727	N7728	476-103	23 37 30.3	26 51 22	-5	9455	21.11	20.96	0.28	9.04	19.60	19.35	0.03	11.18	11.07	0.07	-23.68
A330746	...	...	23 37 32.6	26 53 23	-5	8847	3.46	2.84	0.18	1.22	18.37	17.97	0.08	13.94	13.83	0.06	-20.92
A330747	...	...	23 37 33.6	26 51 24	-2	9799	7.30	6.96	0.85	3.00	20.43	20.15	0.15	15.21	15.09	0.10	-19.65
A330749	...	...	23 37 34.7	26 54 41	-2	9564	4.58	4.11	0.39	1.77	18.88	18.54	0.12	14.64	14.52	0.06	-20.22
A330754	...	476-105	23 37 48.5	27 17 03	-5	8707	18.18	18.01	0.45	7.77	20.27	20.03	0.04	12.13	12.03	0.07	-22.71
A330758	...	...	23 37 51.5	26 54 26	0	10047	7.77	7.44	0.48	3.21	20.01	19.73	0.08	13.75	13.63	0.07	-21.11
A331256	...	...	23 37 57.6	26 49 53	-2	11476	29.71	29.60	2.09	12.76	21.54	21.29	0.08	13.91	13.79	0.06	-20.96
A331259	...	...	23 38 02.5	26 34 43	-2	11548	41.00	40.98	4.66	17.67	22.66	22.42	0.12	15.03	14.92	0.06	-19.83
A331260	...	...	23 38 03.0	26 57 53	-5	10905	10.15	9.87	0.44	4.26	19.95	19.68	0.06	13.19	13.07	0.07	-21.68
A331261	...	...	23 38 06.5	26 30 52	-2	11387	7.13	6.77	0.58	2.92	19.98	19.70	0.11	14.27	14.16	0.06	-20.59
A330761	...	...	23 38 11.8	26 54 01	-2	10198	4.41	3.92	0.45	1.69	18.50	18.16	0.15	14.25	14.13	0.06	-20.61
A331262	...	...	23 38 12.5	26 51 08	-5	9204	5.06	4.63	0.30	2.00	19.34	19.02	0.09	14.00	13.89	0.06	-20.86
A331487	...	...	23 38 14.9	27 23 35	-2	9565	10.19	9.91	0.69	4.27	20.26	20.01	0.09	14.36	14.26	0.06	-20.49
U12733	...	476-107	23 38 16.1	26 33 29	-5	11828	22.06	21.90	0.61	9.44	20.47	20.22	0.04	11.99	11.88	0.07	-22.87
A331489	...	...	23 38 16.2	27 15 41	-2	10307	7.92	7.59	1.00	3.27	21.07	20.81	0.17	15.65	15.55	0.07	-19.20
A331265	...	...	23 38 20.0	26 33 13	-5	8838	9.18	8.85	0.48	3.82	20.01	19.75	0.15	13.38	13.28	0.10	-21.47
A331268	...	...	23 38 22.0	26 58 19	-5	8987	4.87	4.42	0.40	1.91	19.43	19.11	0.12	14.31	14.20	0.06	-20.55
A331491	...	...	23 38 22.1	26 56 05	-5	10667	18.86	18.70	1.32	8.06	21.31	21.05	0.08	13.34	13.22	0.11	-21.52
A330763	...	476-109	23 38 23.6	27 13 55	-5	10127	8.79	8.47	0.27	3.65	19.51	19.25	0.05	12.88	12.78	0.06	-21.97
A331493	...	...	23 38 29.1	26 23 39	-2	9484	23.61	23.46	2.38	10.12	22.24	22.00	0.11	14.73	14.62	0.07	-20.12
A331497	...	...	23 38 50.1	26 55 50	-5	9068	14.00	13.76	1.44	5.93	21.80	21.55	0.12	14.26	14.15	0.10	-20.60
A331272	...	...	23 38 53.6	27 15 15	-2	10033	6.02	5.64	0.70	2.43	20.20	19.92	0.16	14.97	14.87	0.06	-19.88
A331500	...	...	23 39 02.5	27 13 41	-2	9798	11.80	11.55	1.60	4.98	21.30	21.05	0.17	15.43	15.33	0.06	-19.42
A331664	...	...	23 39 25.2	25 59 24	-2	...	39.50	39.50	2.50	17.03	22.85	22.63	0.07	14.73	14.64	0.06	-20.11
U12744	N7735	476-115	23 39 42.0	25 57 00	-5	9596	19.68	19.64	0.35	8.47	20.25	20.03	0.03	11.98	11.89	0.06	-22.86
U12745	N7737	476-118	23 40 15.3	26 46 31	-2	7638	12.05	11.97	0.22	5.16	19.25	19.02	0.03	12.33	12.23	0.05	-22.52

TABLE 3. (continued)

Name	NGC	CGCG	R.A. (1950)	Dec (1950)	T	V_{hel} (km/s)	r_e ($''$)	r_e^c ($''$)	ϵ_r ($''$)	R_e (kpc)	μ_e – (mag/arcsec ²) –	μ_e^c (13)	ϵ_μ (14)	m_I (mag)	m_I^c (mag)	ϵ_m (mag)	M_I (mag)
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)
A331666	...	...	23 40 19.1	27 16 21	-2	...	4.67	4.17	0.32	1.80	19.09	18.77	0.10	14.74	14.64	0.06	-20.11
A331667	...	...	23 40 22.2	25 46 06	-5	...	2.00	1.59	0.19	0.69	18.53	18.12	0.14	15.36	15.28	0.05	-19.46
A330785	...	476-120	23 40 25.2	27 11 40	-5	7566	12.43	12.17	0.27	5.25	19.47	19.23	0.04	12.12	12.03	0.06	-22.72
A331298	N7740	476-122	23 40 44.8	27 05 12	-5	7800	7.55	7.20	0.22	3.10	19.05	18.79	0.05	12.99	12.89	0.06	-21.85
A330788	...	...	23 40 46.0	27 04 55	-2	7566	5.85	5.66	0.18	2.44	18.56	18.30	0.05	13.05	12.95	0.05	-21.79
A330798	...	476-123	23 41 01.1	27 02 04	-2	9031	20.39	20.29	0.78	8.75	19.98	19.75	0.04	12.44	12.34	0.06	-22.41
A331309	...	...	23 41 13.8	27 14 21	-5	...	4.50	3.98	0.19	1.72	18.46	18.13	0.07	13.58	13.48	0.06	-21.27
A331324	...	...	23 43 15.1	26 17 28	-2	...	10.13	10.03	0.61	4.33	19.90	19.67	0.08	14.14	14.05	0.06	-20.70
A330889	...	477-004	23 43 15.9	27 03 42	-2	9143	14.39	14.29	0.65	6.16	20.34	20.11	0.06	13.24	13.14	0.05	-21.60
A331670	...	...	23 43 27.4	26 22 30	0	...	6.86	6.71	0.65	2.89	20.71	20.47	0.13	14.70	14.61	0.07	-20.14
A331333	...	...	23 44 29.2	26 59 07	1	...	62.24	62.24	3.39	26.84	22.65	22.43	0.06	14.02	13.93	0.06	-20.82
A331343	...	...	23 45 14.2	27 14 39	-2	...	18.66	18.57	1.54	8.01	21.55	21.32	0.09	15.31	15.21	0.06	-19.53
A330924	...	477-007	23 45 14.5	27 24 14	-5	6801	4.87	4.64	0.21	2.00	18.41	18.16	0.07	13.14	13.05	0.06	-21.69
A331345	...	...	23 45 27.8	27 16 30	-2	8464	113.1	113.15	5.46	48.80	22.68	22.47	0.04	14.13	14.04	0.05	-20.71
A331416	...	...	23 45 28.2	27 20 26	-2	8188	9.04	8.88	0.79	3.83	19.60	19.37	0.11	14.05	13.96	0.06	-20.79
U12785	...	477-008	23 45 29.8	27 05 50	-2	8381	31.01	31.02	0.99	13.38	20.50	20.28	0.04	12.60	12.51	0.06	-22.24

Notes to Table 3.

A111067; U688; A110044; A111068: within the halo of the galaxy U689.

A110055: bright companion nearby (U700).

U735: faint outer ring.

U878: strong dust lane.

A110246: faint outer ring.

U906: SB0 with outer ring.

A111080: bright companion nearby (U906).

U939: within the halo of the galaxy U938.

U1272: Sa with faint outer ring.

U1277: strong dust lane.

U1308: bright star nearby.

A110848; U1343N; A111121; U1346; A111122; U1345; A111123; U1343S: within the halo of the cD galaxy U1348.

U1406: SB0 with faint outer ring.

U6656: extended low surf. bright. halo.

U6663: SB0 with outer ring.

A211604; A210674; A210679: bright companion nearby (U6704).

A210739: bright companion nearby (U6723).

U6731: SB0 with extended low surf. bright. halo.

U6739: bright companion nearby (U6735).

U8026: bright star nearby.